

Tutorial 7 (Week 7)

Preparatory questions (attempt before the tutorial)

1. Let $A = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 & 0 \\ -1 & -2 & 6 \end{bmatrix}$, $C = [-3 \ 5 \ -1]$, $D = \begin{bmatrix} 3 & 4 \\ 2 & 0 \\ 0 & -7 \\ 1 & -3 \end{bmatrix}$.

Write down the size of each matrix.

2. Let $M = [m_{ij}] = \begin{bmatrix} 6 & 0 & 3 & -5 \\ 0 & 7 & 2 & 4 \\ 1 & 3 & -2 & 0 \end{bmatrix}$. Write down
(a) m_{22} (b) m_{33} (c) m_{14} (d) m_{32} (e) m_{34} .

Tutorial exercises

3. Poole, Exercises 2.4, Question 5:

A coffee merchant sells three blends of coffee. A bag of the house blend has 300 grams of Columbian beans and 200 grams of French roast beans. A bag of the special blend has 200 grams of Columbian beans, 200 grams of Kenyan beans and 100 grams of French roast beans. A bag of the gourmet blend has 100 grams of Columbian beans, 200 grams of Kenyan beans and 200 grams of French roast beans. The merchant has 30 kg of Columbian beans, 15 kg of Kenyan beans and 25 kg of French roast beans, and wishes to use them all up. Determine whether or not he can, and if so, how many bags of each blend he should produce.

4. Poole, Exercises 2.4, Question 6:

Now assume that the house blend has 300 grams of Columbian beans, 50 grams of Kenyan beans and 150 grams of French roast beans, the special blend has 200 grams of Columbian, 200 grams of Kenyan and 100 grams of French roast and the gourmet blend has 100 grams of Columbian, 350 grams of Kenyan and 50 grams of French roast. The merchant has 30 kg of Columbian beans, 15 kg of Kenyan and 15 kg of French roast available. The respective profits per bag are \$0.50, \$1.50 and \$2.00 for the house, special and gourmet blends.

The merchant wants to package *all* the beans *and* maximise his profit. How many bags of each type should he produce, and what is the resulting profit?

5. Poole, Exercises 2.4, Questions 8 and 12:

Balance the chemical equation for each of the following reactions:

- (a) $\text{CO}_2 + \text{H}_2\text{O} \longrightarrow \text{C}_6\text{H}_{12}\text{O}_6 + \text{O}_2$
(b) $\text{HClO}_4 + \text{P}_4\text{O}_{10} \longrightarrow \text{H}_3\text{PO}_4 + \text{Cl}_2\text{O}_7$

6. Poole, Exercises 2.4, Question 32:

The sum of the ages of Annie, Bert and Chris is 60. Annie is older than Bert by the same number of years as Bert is older than Chris. When Bert is as old as Annie is now, Annie will be three times as old as Chris is now. What are their ages?

7. Let $A = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 \\ -1 & -2 \end{bmatrix}$, $C = \begin{bmatrix} 0 & 2 \\ 4 & -3 \end{bmatrix}$, $D = \begin{bmatrix} 10 & 0 \\ 0 & 5 \end{bmatrix}$.

Find each of the following.

- (a) $A + B$ (e) $2A$ (i) AB (m) $A(BC)$ (q) B^2
(b) $A - B$ (f) $-B$ (j) BA (n) $(AB)C$ (r) A^2B^2
(c) $B - C$ (g) $3C$ (k) CD (o) $ABCD$
(d) $D + C$ (h) $\frac{1}{5}D$ (l) BC (p) A^2

8. Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ 1 & -1 \\ -1 & 2 \end{bmatrix}$, $C = [3 \ 4 \ 2]$, $D = \begin{bmatrix} -1 \\ -1 \\ 5 \end{bmatrix}$.

Find any of the following that exist:

$AB, AC, AD, BA, BC, BD, CA, CB, CD, DA, DB, DC$.

9. Find a 2×2 matrix M such that $M^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ but every entry of M is nonzero.

Further exercises

In addition to these exercises, the following exercises from the textbook – *Linear Algebra: A Modern Introduction* by David Poole – should be attempted:

Exercises 2.4: 1, 3, 7, 9, 11, 31, 33, 39.

Exercises 3.1: 1, 3, 5, 7, 9, 11, 15.

Answers to selected exercises

1. 2×2 ; 2×3 ; 1×3 ; 4×2 .
2. (a) 7 (b) -2 (c) -5 (d) 3 (e) 0
3. 65 bags of house blend, 30 bags of special blend and 45 bags of gourmet blend.
4. 60 bags of house blend, 60 bags of special blend, no bags of gourmet blend.
Maximum profit = \$120.
5. (a) $6 \text{CO}_2 + 6 \text{H}_2\text{O} \longrightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{O}_2$
(b) $12 \text{HClO}_4 + \text{P}_4\text{O}_{10} \longrightarrow 4 \text{H}_3\text{PO}_4 + 6 \text{Cl}_2\text{O}_7$
6. Chris is 12, Bert is 20 and Annie is 28.

$$\begin{array}{ll}
 \text{7. (a) } A + B = \begin{bmatrix} 4 & 2 \\ -2 & 1 \end{bmatrix} & \text{(j) } BA = \begin{bmatrix} -1 & 6 \\ 1 & -4 \end{bmatrix} \\
 \text{(b) } A - B = \begin{bmatrix} -2 & -6 \\ 0 & 5 \end{bmatrix} & \text{(k) } CD = \begin{bmatrix} 0 & 10 \\ 40 & -15 \end{bmatrix} \\
 \text{(c) } B - C = \begin{bmatrix} 3 & 2 \\ -5 & 1 \end{bmatrix} & \text{(l) } BC = \begin{bmatrix} 16 & -6 \\ -8 & 4 \end{bmatrix} \\
 \text{(d) } D + C = \begin{bmatrix} 10 & 2 \\ 4 & 2 \end{bmatrix} & \text{(m) } A(BC) = \begin{bmatrix} 32 & -14 \\ -40 & 18 \end{bmatrix} \\
 \text{(e) } 2A = \begin{bmatrix} 2 & -4 \\ -2 & 6 \end{bmatrix} & \text{(n) } (AB)C = \begin{bmatrix} 32 & -14 \\ -40 & 18 \end{bmatrix} \\
 \text{(f) } -B = \begin{bmatrix} -3 & -4 \\ 1 & 2 \end{bmatrix} & \text{(o) } ABCD = \begin{bmatrix} 320 & -70 \\ -400 & 90 \end{bmatrix} \\
 \text{(g) } 3C = \begin{bmatrix} 0 & 6 \\ 12 & -9 \end{bmatrix} & \text{(p) } A^2 = \begin{bmatrix} 3 & -8 \\ -4 & 11 \end{bmatrix} \\
 \text{(h) } \frac{1}{5}D = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} & \text{(q) } B^2 = \begin{bmatrix} 5 & 4 \\ -1 & 0 \end{bmatrix} \\
 \text{(i) } AB = \begin{bmatrix} 5 & 8 \\ -6 & -10 \end{bmatrix} & \text{(r) } A^2B^2 = \begin{bmatrix} 23 & 12 \\ -31 & -16 \end{bmatrix}
 \end{array}$$

$$\text{8. } AB = \begin{bmatrix} -1 & 5 \\ 1 & 3 \end{bmatrix}, \quad AD = \begin{bmatrix} 12 \\ 0 \end{bmatrix}, \quad BA = \begin{bmatrix} 3 & 2 & 1 \\ -2 & 0 & 2 \\ 5 & 2 & -1 \end{bmatrix}, \quad CB = [2 \ 3], \\
 CD = [3], \quad DC = \begin{bmatrix} -3 & -4 & -2 \\ -3 & -4 & -2 \\ 15 & 20 & 10 \end{bmatrix}.$$

The other products are not defined.