

The short solutions suffice except for the following questions.

4. (ii) The inverse of $\begin{bmatrix} 6 & 2 \\ 3 & 1 \end{bmatrix}$ does not exist because its determinant is $6(1) - 2(3) = 0$.

$$(iv) \left[\begin{array}{ccc|ccc} 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & 0 \end{array} \right]$$

$$(v) \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & -1 & -2 & 0 & 1 & 0 \\ -1 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & -1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & -1 & 0 \\ 0 & 0 & -1 & 1 & 1 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 2 & 1 & 1 \\ 0 & 1 & 0 & 2 & 1 & 2 \\ 0 & 0 & 1 & -1 & -1 & -1 \end{array} \right]$$

$$(vi) \left[\begin{array}{ccc|ccc} 2 & 4 & 6 & 1 & 0 & 0 \\ 7 & 11 & 6 & 0 & 1 & 0 \\ -6 & -6 & 12 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1/2 & 0 & 0 \\ 0 & -3 & -15 & -7/2 & 1 & 0 \\ 0 & 6 & 30 & 3 & 0 & 1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1/2 & 0 & 0 \\ 0 & -3 & -15 & -7/2 & 1 & 0 \\ 0 & 0 & 0 & * & * & * \end{array} \right],$$

so the matrix is not invertible.

$$(vii) \left[\begin{array}{ccc|ccc} -4 & 3 & 3 & 1 & 0 & 0 \\ 8 & 7 & 3 & 0 & 1 & 0 \\ 4 & 3 & 3 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} -4 & 3 & 3 & 1 & 0 & 0 \\ 0 & 13 & 9 & 2 & 1 & 0 \\ 0 & 6 & 6 & 1 & 0 & 1 \end{array} \right]$$

$$\sim \left[\begin{array}{ccc|ccc} 1 & -3/4 & -3/4 & -1/4 & 0 & 0 \\ 0 & 1 & 1 & 1/6 & 0 & 1/6 \\ 0 & 0 & -4 & -1/6 & 1 & -13/6 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -1/8 & 0 & 1/8 \\ 0 & 1 & 0 & 1/8 & 1/4 & -3/8 \\ 0 & 0 & 1 & 1/24 & -1/4 & 13/24 \end{array} \right]$$

5. By the formula for 2×2 matrices, the inverse of $\begin{bmatrix} 5 & -3 \\ 7 & -4 \end{bmatrix}$ is $\begin{bmatrix} -4 & 3 \\ -7 & 5 \end{bmatrix}$, so that

$$\begin{bmatrix} x & y \\ z & w \end{bmatrix} = \begin{bmatrix} -4 & 3 \\ -7 & 5 \end{bmatrix} \begin{bmatrix} 11 & 4 \\ 15 & 5 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ -2 & -3 \end{bmatrix}.$$

7. Observe that

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 3 & 0 & 1 & 0 \\ 3 & 4 & 3 & 0 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -2 & 1 & 0 \\ 0 & 1 & 0 & -3 & 0 & 1 \end{array} \right] \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 6 & -1 & -1 \\ 0 & 1 & 0 & -3 & 0 & 1 \\ 0 & 0 & 1 & -2 & 1 & 0 \end{array} \right],$$

so the inverse of $\begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 3 \\ 3 & 4 & 3 \end{bmatrix}$ is $\begin{bmatrix} 6 & -1 & -1 \\ -3 & 0 & 1 \\ -2 & 1 & 0 \end{bmatrix}$. Observe also that

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 3 \\ 3 & 4 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix},$$

so that

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 & -1 & -1 \\ -3 & 0 & 1 \\ -2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 11 \\ -5 \\ -4 \end{bmatrix}.$$

9. (i) This is false. For example take

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Then

$$(ABC)^{-1} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix},$$

yet

$$A^{-1}B^{-1}C^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

(ii) This is true, since $(ABA)^{-1} = A^{-1}(AB)^{-1} = A^{-1}B^{-1}A^{-1}$.

(iii) This is true. By uniqueness of inverses, since $A^{-1}A = AA^{-1} = I$, we have immediately that $(A^{-1})^{-1} = A$.

(iv) This is true. Observe that

$$(-A)(-A^{-1}) = (-1)(-1)AA^{-1} = AA^{-1} = I$$

and

$$(-A^{-1})(-A) = (-1)(-1)A^{-1}A = A^{-1}A = I,$$

so that, by uniqueness of inverses, $(-A)^{-1} = -A^{-1}$, yielding

$$-(-A)^{-1} = -(-A^{-1}) = A^{-1}.$$

(v) This is true, since $C^{-1}(ABC^{-1})^{-1}AB = C^{-1}(C^{-1})^{-1}B^{-1}A^{-1}AB = I$.

(vi) This is false even for 1×1 matrices, since $(A+B)^{-1}$ may not exist. For example, take $A = 1$ and $B = -1$, so that $A+B = 0$ has no inverse. Even when $(A+B)^{-1}$ exists, the statement is typically false. For example, take $A = B = 1$, so that $(A+B)^{-1} = 1/2 \neq 2 = A^{-1} + B^{-1}$.

(vii) This is true, since $A^{-1}(I+A)A = A^{-1}IA + A^{-1}AA = I + A = A + I$.

(viii) This is true, since $(A+I)(A^{-1}-I) = AA^{-1} - A + A^{-1} - I = A^{-1} - A$.

(ix) This is true, since

$$\begin{aligned} A^2 - 2A + I = 0 & \implies 2A - A^2 = I \\ & \implies A(2I - A) = (2I - A)A = I \\ & \implies A^{-1} = 2I - A. \end{aligned}$$

(x)* This is false. For example, take $A = \begin{bmatrix} 2 & 1 \\ -1 & 0 \end{bmatrix} \neq I$, yet

$$A^2 - 2A + I = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} - \begin{bmatrix} 4 & 2 \\ -2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 0.$$

10.* Observe that

$$\begin{bmatrix} 1 & -2 & 3 \\ -3 & 1 & 2 \\ -3 & -4 & \lambda \end{bmatrix} \sim \begin{bmatrix} 1 & -2 & 3 \\ 0 & -5 & 11 \\ 0 & -10 & \lambda + 9 \end{bmatrix} \sim \begin{bmatrix} 1 & -2 & 3 \\ 0 & -5 & 11 \\ 0 & 0 & \lambda - 13 \end{bmatrix},$$

so that the method of finding the inverse by row reduction fails if and only if $\lambda = 13$.

11. Observe that $(5M)^{-1} = \begin{bmatrix} 5 & 6 \\ 5 & 5 \end{bmatrix}$ so that $5M = \begin{bmatrix} 5 & 6 \\ 5 & 5 \end{bmatrix}^{-1} = -\frac{1}{5} \begin{bmatrix} 5 & -6 \\ -5 & 5 \end{bmatrix}$, yielding

$$M = -\frac{1}{25} \begin{bmatrix} 5 & -6 \\ -5 & 5 \end{bmatrix} = \begin{bmatrix} -1/5 & 6/25 \\ 1/5 & -1/5 \end{bmatrix}.$$

12. Observe that

$$\begin{bmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 2 & 3 & 1 & | & 0 & 1 & 0 \\ 3 & 1 & 2 & | & 0 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 3 & | & 1 & 0 & 0 \\ 0 & -1 & -5 & | & -2 & 1 & 0 \\ 0 & -5 & -7 & | & -3 & 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -7 & | & -3 & 2 & 0 \\ 0 & 1 & 5 & | & 2 & -1 & 0 \\ 0 & 0 & 18 & | & 7 & -5 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & | & -5/18 & 1/18 & 7/18 \\ 0 & 1 & 0 & | & 1/18 & 7/18 & -5/18 \\ 0 & 0 & 1 & | & 7/18 & -5/18 & 1/18 \end{bmatrix}$$

so the inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix}$ is $\frac{1}{18} \begin{bmatrix} -5 & 1 & 7 \\ 1 & 7 & -5 \\ 7 & -5 & 1 \end{bmatrix}$. Observe also that

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix},$$

so that

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{18} \begin{bmatrix} -5 & 1 & 7 \\ 1 & 7 & -5 \\ 7 & -5 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \frac{1}{18} \begin{bmatrix} -5a + b + 7c \\ a + 7b - 5c \\ 7a - 5b + c \end{bmatrix}.$$

13. If any of the diagonal entries is zero, then the matrix has a row of zeros so is not invertible. If all of the diagonal entries are nonzero then

$$\left[\begin{array}{cccc|cccc} d_1 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_n & 0 & 0 & \cdots & 1 \end{array} \right] \sim \left[\begin{array}{cccc|cccc} 1 & 0 & \cdots & 0 & d_1^{-1} & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 & 0 & d_2^{-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 & 0 & 0 & \cdots & d_n^{-1} \end{array} \right]$$

so that the inverse exists and is the diagonal matrix with reciprocals down the diagonal.

14.* If $n = 1$ then $I - J = 1 - 1 = 0$ which is not invertible. Suppose $n \geq 2$. Then $J^2 = nJ$, so that

$$(I - J)\left(I - \frac{1}{n-1}J\right) = I - \frac{1}{n-1}J - J + \frac{1}{n-1}J^2 = I - \frac{n}{n-1}J + \frac{n}{n-1}J = I,$$

and similarly $\left(I - \frac{1}{n-1}J\right)(I - J) = I$, so that $(I - J)^{-1} = I - \frac{1}{n-1}J$.

15.* (i) Observe that $\begin{bmatrix} 2-\lambda & 0 \\ 0 & -3-\lambda \end{bmatrix} \sim \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ if and only if $\lambda \neq 2$ and $\lambda \neq -3$,

so that $A - \lambda I$ is not invertible if and only if $\lambda = 2$ or $\lambda = -3$.

(ii) Observe that

$$\begin{bmatrix} 1-\lambda & 2 \\ -1 & 4-\lambda \end{bmatrix} \sim \begin{bmatrix} 1 & \lambda-4 \\ 1-\lambda & 2 \end{bmatrix} \sim \begin{bmatrix} 1 & \lambda-4 \\ 0 & \lambda^2-5\lambda+6 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

if and only if $\lambda^2 - 5\lambda + 6 = (\lambda - 2)(\lambda - 3) \neq 0$. Hence $A - \lambda I$ is not invertible if and only if $(\lambda - 2)(\lambda - 3) = 0$, that is, $\lambda = 2$ or $\lambda = 3$.

(iii) Observe that

$$\begin{aligned} & \begin{bmatrix} -3-\lambda & 0 & 2 \\ -4 & -1-\lambda & 4 \\ -4 & -4 & 7-\lambda \end{bmatrix} \sim \begin{bmatrix} -4 & -4 & 7-\lambda \\ 0 & 3-\lambda & \lambda-3 \\ -3-\lambda & 0 & 2 \end{bmatrix} \\ & \sim \begin{bmatrix} 1 & 1 & (\lambda-7)/4 \\ 0 & 3-\lambda & \lambda-3 \\ 0 & \lambda+3 & (\lambda^2-4\lambda-13)/4 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & (\lambda-7)/4 \\ 0 & 3-\lambda & \lambda-3 \\ 0 & 6 & (\lambda^2-25)/4 \end{bmatrix} \\ & \sim \begin{bmatrix} 1 & 1 & (\lambda-7)/4 \\ 0 & 1 & (\lambda^2-25)/24 \\ 0 & 3-\lambda & \lambda-3 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & (\lambda-7)/4 \\ 0 & 1 & (\lambda^2-25)/24 \\ 0 & 0 & (\lambda-3)(\lambda-1)(\lambda+1)/24 \end{bmatrix}, \end{aligned}$$

which can be row reduced to the identity matrix if and only if

$$(\lambda - 3)(\lambda - 1)(\lambda + 1) \neq 0.$$

Hence $A - \lambda I$ is not invertible if and only if

$$(\lambda - 3)(\lambda - 1)(\lambda + 1) = 0,$$

that is, $\lambda = 3, 1$ or -1 .