

ASYMPTOTIC ANALYSIS OF RICCI FLOW ON \mathbb{R}^{n+1} WITH TYPE-IIB SINGULARITIES

HAOTIAN WU

ABSTRACT. In this paper, we study the precise asymptotics of Type-IIb solutions to Ricci flow on \mathbb{R}^{n+1} . In each dimension $n + 1 \geq 3$ and for each real number $\lambda > 0$, we construct a family of complete rotationally symmetric solutions to Ricci flow on \mathbb{R}^{n+1} that form in infinite time Type-IIb singularities with the curvature blow-up rate $t^{\lambda-1}$. Near the origin, the blow-ups of such a solution converge uniformly to the Bryant soliton; near spatial infinity, the solution is asymptotically flat at a precise rate depending on λ .

1. INTRODUCTION

A one-parameter family of $(n+1)$ -dimensional complete smooth Riemannian manifolds $(M, g(t))_{t_0 \leq t < t_1}$ is said to evolve by Hamilton's Ricci flow [19], starting from an initial metric g_0 , if $g(t)$ satisfies the equation

$$(1.1) \quad \partial_t g = -2 \operatorname{Ric}(g), \quad g(t_0) = g_0,$$

where $\operatorname{Ric}(g)$ is the Ricci curvature of the metric.

Let $(M, g(t))$ be a solution to Ricci flow that exists up to a maximal time $T \leq \infty$. If $T < \infty$, then we say the Ricci flow has a finite-time singularity of

- Type-I if $\sup_{M \times [t_0, T)} (T - t) |\operatorname{Rm}(\cdot, t)| < \infty$,
- Type-IIa if $\sup_{M \times [t_0, T)} (T - t) |\operatorname{Rm}(\cdot, t)| = \infty$.

If $T = \infty$, then the infinite time singularity of this immortal Ricci flow is said to be

- Type-III if $\sup_{M \times [t_0, \infty)} t |\operatorname{Rm}(\cdot, t)| < \infty$,
- Type-IIb if $\sup_{M \times [t_0, \infty)} t |\operatorname{Rm}(\cdot, t)| = \infty$.

If a Ricci flow solution encounters a Type-IIb (or Type-III) singularity, then we also call it a Type-IIb (or Type-III) solution to Ricci flow. Analogous classifications hold for solutions to mean curvature flow (MCF) with $|h(\cdot, t)|^2$,

Date: November 18, 2019.

2010 Mathematics Subject Classification. 53C44 (primary), 35K59 (secondary).

Key words and phrases. Ricci flow; noncompact manifolds; Type-IIb singularity; precise asymptotics.

the second fundamental form of a hypersurface moving by MCF, replacing $|\text{Rm}(\cdot, t)|$ in the above definitions.

Naturally, we have the following questions (cf. [11, Problem 8.6]): What can be said about the specific blow-up rates of Ricci flow solutions with a Type-IIa or Type-IIb singularity? What about the asymptotic properties Ricci flow solutions of each type near the singular time T ?

In real dimension two ($n+1 = 2$), by the work of Hamilton [20] and Chow [9], Ricci flow on S^2 always encounters a Type-I singularity modelled by the round sphere. On the other hand, Daskalopoulos and Hamilton [14] have showed that Ricci flow on \mathbb{R}^2 starting from a metric of finite area forms a Type-IIa singularity at the rate $(T-t)^{-2}$. The precise description of the extinction profile of such a solution were later given in [13] and [15]: the solution is modelled by a cigar soliton in an inner region, and has a logarithmic cusp in an outer region.

In real dimension three or higher ($n+1 \geq 3$), Hamilton's seminal work [19] says that Ricci flow of a closed three-manifold with positive Ricci curvature forms a Type-I singularity and shrinks to a round point. This result was later generalised to higher dimensions under other curvature assumptions, e.g., the 2-positive curvature operator by Böhm and Wilking [4]. These Type-I singularities are global in the sense that the volume of the manifold at the singular time T is zero. In comparison, there exist local singularities that form on compact subsets of a manifold and the volume of the manifold remains positive at the singular time T . For example, Type-I non-degenerate neckpinches modelled by the round cylinder have been rigorously constructed by Angenent and Knopf [1].

In real dimensions $n+1 \geq 3$, Type-IIa singularity was first proved to exist in Ricci flow on S^{n+1} by Gu and Zhu [18]. Concerning the geometric details of such a solution, Garfinkle and Isenberg [17] gave numerical evidence that a degenerate neckpinch in Ricci flow on S^3 is a Type-IIa singularity modelled by the rotationally symmetric Bryant soliton, which was first constructed by Bryant [7] and has been proven by Brendle to be the unique complete non-flat steady gradient Ricci soliton in dimension three under a non-collapsing assumption [5]; see also Brendle's generalisation to higher dimensions [6]. In [2], Angenent, Isenberg and Knopf have constructed on S^{n+1} Ricci flow with Type-IIa singularities modelled on the Bryant soliton with curvature blow-up rate $(T-t)^{-2+2/k}$ for each integer $k \geq 3$. In contrast, Type-IIa singularities to Ricci flow on \mathbb{R}^n with curvature blow-up rates $(T-t)^{-\lambda+1}$ for any real number $\lambda \geq 1$ have been constructed by the author in [29]. There are also corresponding results on Type-IIa singularities in MCF by the author and his collaborators [21, 22].

There are several recent results on Ricci flow with Type-IIa singularities. Appleton [3] has showed that Ricci flow on a noncompact four-manifold can develop Type-IIa singularities modelled on the Eguchi-Hanson space. Di Giovanni [16] has proved that asymptotically cylindrical Ricci flow on \mathbb{R}^{n+1} without minimal sphere forms a Type-IIa singularity modelled on the

Bryant soliton after suitable dilations. Stolarski [27] has constructed on certain product manifolds Ricci flows that form Type-IIa singularities with curvature blow-up rates given by arbitrarily large powers of $(T - t)$. If we specialise the Ricci flow to Kähler manifolds, then Li, Tian and Zhu have given the first examples of Type-IIa singularities for the Kähler-Ricci flow [24].

Concerning the Type-IIb singularities in Ricci flow, the simplest example on compact manifolds is a nonflat Ricci-flat Kähler metric on a K3 surface, whose existence follows from Yau’s resolution [30] of the Calabi conjecture; note that this solution is static under Ricci flow. Further results on Kähler-Ricci flows with Type-IIb singularities have been obtained by Tosatti and Zhang [28]. It has been conjectured [10] that Ricci flow on closed 3-manifolds never form Type-IIb singularity. On a noncompact manifold, the Bryant soliton is trivially a Ricci flow solution with Type-IIb singularity. In this paper, we are interested in finding *non-Kähler, non-Bryant-soliton*, solutions to Ricci flow with Type-IIb singularities on a complete noncompact manifold.

Throughout this paper, we use C_k ($k \in \mathbb{N}$) to denote a positive constant that depends at most on n or λ , and may change from line to line. The expression “ $f \lesssim g$ ” means $f \leq C_k g$ for some constant C_k ; “ $A \sim B$ ” if and only if $A \lesssim B$ and $B \lesssim A$.

Our main result is the following.

Theorem 1.1. *In each dimension $n + 1 \geq 3$, for each real number $\lambda > 0$, there exists a family \mathcal{G} of complete rotationally symmetric metrics, none of which is the Bryant soliton, on \mathbb{R}^{n+1} such that Ricci flow starting at each $g_0 \in \mathcal{G}$ has a unique solution $g(t)$ for $t \in [t_0, \infty)$. The solution $g(t)$ has the following asymptotic properties as $t \nearrow \infty$.*

- (1) *The singularity is Type-IIb with*

$$\sup_{\mathbb{R}^{n+1}} |\text{Rm}(\cdot, t)| \sim t^{\lambda-1}$$

attained at the origin of \mathbb{R}^{n+1} .

- (2) *If we rescale the solution so that the distance from the origin rescales at the rate $t^{-(1-\lambda)/2}$, then the metric converges uniformly on intervals of order $t^{(1-\lambda)/2}$ to the Bryant soliton.*
- (3) *Near spatial infinity, the metric is asymptotically flat, i.e., $|\text{Rm}(\cdot, t)| \rightarrow 0$, for all $t \geq t_0$. More precisely, the sectional curvatures K and L as defined in (2.3) satisfy*

$$K \sim \psi^{-2(1+\lambda)}, \quad L \sim \psi^{-2}$$

as $\psi \nearrow \infty$, for all $t \geq t_0$.

In particular, the solution exhibits the asymptotic behaviour of the formal solution described in Section 3.

To the author's knowledge, Theorem 1.1 gives the first examples of non-Kähler Type-IIb solutions to Ricci flow on \mathbb{R}^{n+1} for $n \geq 2$ that are not the Bryant soliton. These solutions (and also the Bryant soliton) show that the exponent $(\lambda - 1)$ of the Type-IIb blow-up rate $t^{\lambda-1}$ belongs to a continuum $(-1, \infty)$. We note that a similar phenomenon has been observed for Type-IIa singularities in Ricci flows on \mathbb{R}^{n+1} [29]. As $t \nearrow \infty$, the Ricci flow solutions constructed in Theorem 1.1 converge uniformly to a non-Euclidean flat metric (cf. Remark 3.2) metric if $\lambda \in (0, 1)$ and otherwise if $\lambda \geq 1$. As previously mentioned, the Bryant soliton is a Type-IIb solution whose curvature blow-up rate is $t^{\lambda-1}$ with $\lambda = 1$. So we may ask whether or not the Bryant soliton appears as a "phase change" among Type-IIb solutions to Ricci flow when the parameter λ varies across the "critical value" $\lambda = 1$. Lastly, one may compare Theorem 1.1 for Ricci flow with the construction of Type-IIb solutions to MCF in [23].

The proof of Theorem 1.1 uses matched asymptotic analysis and barrier arguments for nonlinear PDE. The same strategy has been implemented for Ricci flow or mean curvature flow with Type-IIa singularities in [21, 22, 29], and Type-IIb MCF solutions in [23]. In Section 2, we recall the set-up for rotationally symmetric Ricci flow on \mathbb{R}^{n+1} and collect some basic facts. In Section 3, we derive approximate (formal) solutions using the method of formal matched asymptotics. In Section 4, we use these approximate solutions to construct the corresponding supersolutions and subsolutions to the rescaled PDE. The supersolutions and subsolutions are ordered and patched together in Section 5 to create barriers to the rescaled PDE; a comparison principle for the subsolutions and supersolutions is also proved there. In Section 6, we complete the proof of Theorem 1.1.

Acknowledgements. The author is supported by ARC grant DE180101348.

2. PRELIMINARIES

Let \mathcal{O} denote the origin of \mathbb{R}^{n+1} ($n \geq 2$). We identify $\mathbb{R}^{n+1} \setminus \mathcal{O}$ with $(0, \infty) \times S^n$ and equip it with the time-dependent warped product metric

$$g = \varphi^2(x, t)dx^2 + \psi^2(x, t)g_{\text{sph}},$$

where $x \in (0, \infty)$ and g_{sph} is the metric of constant sectional curvature one on S^n .

We recall some basic facts about such a metric, cf. [2, Section 2]. The distance s to the origin is

$$s(x, t) := \int_0^x \varphi(y, t)dy.$$

In the s -coordinate, the metric becomes

$$(2.1) \quad g = ds^2 + \psi^2(s, t)g_{\text{sph}}.$$

If we extend the metric g to a complete smooth metric, still denoted by g , on \mathbb{R}^{n+1} , then ψ necessarily satisfy the boundary conditions

$$\lim_{x \searrow 0} \psi = 0 \quad \text{and} \quad \lim_{x \searrow 0} \psi_s = 1.$$

In this paper, we use the notation $\partial_t|_x$ for taking the time derivative while keeping the quantity “.” fixed. Then

$$[\partial_t|_x, \partial_s] = -n \frac{\psi_{ss}}{\psi} \partial_s.$$

In the s -coordinate, the Ricci flow system (1.1) is reduced to the following parabolic PDE for ψ ,

$$(2.2) \quad \partial_t|_x \psi = \psi_{ss} - (n-1) \frac{1 - \psi_s^2}{\psi}.$$

The function φ , which is suppressed in the s -coordinate, evolves under Ricci flow by

$$\partial_t|_x \log \varphi = n \frac{\psi_{ss}}{\psi}.$$

Let K denote the sectional curvature of a two-plane with one radial and one spherical direction and L the sectional curvature of a two-plane tangential to the sphere $\{x\} \times S^n$. Then

$$(2.3) \quad K = -\frac{\psi_{ss}}{\psi}, \quad L = \frac{1 - \psi_s^2}{\psi^2}.$$

In particular, $|\text{Rm}|^2 = 2nK^2 + n(n-1)L^2$.

Since the metric g is smooth and $\lim_{x \searrow 0} \psi_s = 1$, we must have $\psi_s > 0$ in a neighbourhood of the origin \mathcal{O} . So we can use ψ as a new coordinate near the origin to write

$$(2.4) \quad g = z(\psi, t)^{-1} d\psi^2 + \psi^2 g_{\text{sph}},$$

where $z(\psi, t) := \psi_s^2$. Then the sectional curvatures are rewritten as

$$K = -\frac{z\psi}{2\psi}, \quad L = \frac{1-z}{\psi^2}.$$

Under Ricci flow, the metric (2.4) evolves by

$$(2.5) \quad \partial_t|_\psi z = \mathcal{E}_\psi[z],$$

where \mathcal{E}_ψ is the purely local quasilinear operator

$$\mathcal{E}_\psi[z] := zz_{\psi\psi} - \frac{1}{2}z_\psi^2 + (n-1-z)\frac{z\psi}{\psi} + 2(n-1)\frac{(1-z)z}{\psi^2}.$$

We can split \mathcal{E}_ψ into a linear part and a quadratic part:

$$\mathcal{E}_\psi[z] = \mathcal{L}_\psi[z] + \mathcal{Q}_\psi[z],$$

where

$$(2.6) \quad \mathcal{L}_\psi[z] := (n-1) \left(\frac{z_\psi}{\psi} + 2 \frac{z}{\psi^2} \right),$$

$$(2.7) \quad \mathcal{Q}_\psi[z] := zz_\psi - \frac{1}{2} z_\psi^2 - \frac{zz_\psi}{\psi} - 2(n-1) \frac{z^2}{\psi^2}.$$

The quadratic part defines a symmetric bilinear operator

$$(2.8) \quad \hat{\mathcal{Q}}_\psi[z_1, z_2] := \frac{1}{2} [z_1(z_2)_\psi + z_2(z_1)_\psi - (z_1)_\psi(z_2)_\psi]$$

$$(2.9) \quad - \frac{z_1(z_2)_\psi + z_2(z_1)_\psi}{2\psi} - 2(n-1) \frac{z_1 z_2}{\psi^2}.$$

In particular, $\mathcal{Q}_\psi[z] = \hat{\mathcal{Q}}_\psi[z, z]$.

3. FORMAL SOLUTIONS

The basic idea behind the construction of the formal solutions is to analyse equation (2.5) under various rescalings of ψ and find approximate solutions to the rescaled PDEs. We introduce the following rescaled variables

$$\begin{aligned} \tau &:= \log t, \\ \sigma &:= \frac{s}{\sqrt{2(n-1)t}}, \\ u &:= \frac{\psi}{\sqrt{2(n-1)t}}. \end{aligned}$$

Since we are interested in the asymptotic behaviour of the solution when $t \nearrow \infty$, we can assume $t_0 \geq 1$, and so $\tau_0 = \log t_0 \geq 0$.

In the (u, τ) -coordinates, equation (2.5) becomes

$$(3.1) \quad \partial_\tau|_u z = \frac{1}{2(n-1)} \mathcal{E}_u[z] + \frac{1}{2} uz_u,$$

where $\tau \in [\log t_0, \infty)$, $u \in (0, \infty)$ and $z \in (0, 1]$. In particular, $z \in (0, 1]$ under Ricci flow, which is proved in Lemma 6.2. We seek solutions of equation (3.1) subject to the boundary condition $z(0, \tau) = 1$ and the asymptotic condition $\lim_{u \nearrow \infty} z(u, \tau) = 0$ for all $\tau \geq \tau_0$. In particular, the asymptotic condition that $z \searrow 0$ as $u \nearrow \infty$ is compatible with the consideration of asymptotically flat Riemannian manifolds whose metrics $ds^2 + \psi(s)^2 g_{\text{sph}}$ are defined by $\psi(s)$ with sublinear growth in s , cf. Section 3.4.

3.1. Formal solution in the exterior region. The exterior region is expected to be a time-dependent subset in which $u \in (0, \infty)$ and $z \in (0, 1)$. Motivated by the asymptotic condition $\lim_{u \nearrow \infty} z(u, \tau) = 0$ for all $\tau \geq \tau_0$, we adopt the following ansatz

$$z = \sum_{m=1}^{\infty} e^{-m\lambda\tau} Z_m(u),$$

where $\lambda > 0$ is a parameter to be specified.

We substitute this ansatz into equation (3.1) and split $\mathcal{E}_u[z]$ into the linear and quadratic parts as given in (2.6) and (2.7), respectively. By comparing the coefficients of $e^{-m\lambda\tau}$ in the resulting equation, we see each Z_m must satisfy the ODE

$$(3.2) \quad \frac{1}{2} (u^{-1} + u) \frac{dZ_m}{du} + (u^{-2} + m\lambda) Z_m = -\frac{1}{2(n-1)} \sum_{i=1}^{m-1} \hat{Q}_u [Z_i, Z_{m-i}].$$

When $m = 1$, Z_1 satisfies the linear homogeneous equation

$$(3.3) \quad \frac{1}{2} (u^{-1} + u) \frac{dZ_1}{du} + (u^{-2} + \lambda) Z_1 = 0,$$

whose general solution is

$$Z_1(u) = cu^{-2} (1 + u^2)^{1-\lambda}$$

for an arbitrary constant $c \neq 0$.

When $m = 2$, equation (3.2) becomes

$$(3.4) \quad \frac{1}{2} (u^{-1} + u) \frac{dZ_2}{du} + (u^{-2} + 2\lambda) Z_2 = -\frac{1}{2(n-1)} Q_u [Z_1],$$

where

$$Q_u [Z_1] = 2c^2 u^{-6} (1 + u^2)^{-2\lambda} (4 - n(1 + u^2)^2 + u^4(1 + \lambda)^2 + 2u^2(3 + \lambda)).$$

The general solution of equation (3.4) is

$$Z_2(u) = \frac{u^{-2} (1 + u^2)^{1-2\lambda}}{n-1} f(u),$$

where

$$(3.5) \quad f(u) := C_1 - 2c^2 \left(\frac{n-4}{u^2} + \frac{1-\lambda^2}{1+u^2} \right) - 2c^2(\lambda-1) \log \left(\frac{u^2}{1+u^2} \right)$$

for some arbitrary constant C_1 .

Let us now analyse the asymptotics of $e^{-\lambda\tau} Z_1(u) + e^{-2\lambda\tau} Z_2(u)$ as $u \searrow 0$ and $u \nearrow \infty$, respectively. It is straightforward to see that

$$\frac{e^{-2\lambda\tau} Z_2(u)}{e^{-\lambda\tau} Z_1(u)} = \frac{e^{-\lambda\tau}}{c(n-1)} (1 + u^2)^{-\lambda} f(u),$$

where $f(u)$ as defined in (3.5) has the following asymptotics

$$f(u) \sim \begin{cases} c^2 u^{-2} + O(c^2 \log(u^2)), & u \searrow 0, \\ C_1 + c^2 u^{-2}, & u \nearrow \infty. \end{cases}$$

Therefore, we obtain

$$\frac{e^{-2\lambda\tau} Z_2(u)}{e^{-\lambda\tau} Z_1(u)} = \begin{cases} ce^{-\lambda\tau} (u^{-2} + O(\log u)), & u \searrow 0, \\ e^{-\lambda\tau} u^{-2\lambda} (C_1 + c^2 u^{-2}), & u \nearrow \infty. \end{cases}$$

Consequently, we always have

$$\lim_{u \nearrow \infty} \left| \frac{e^{-2\lambda\tau} Z_2(u)}{e^{-\lambda\tau} Z_1(u)} \right| = 0$$

for all $\tau \geq \tau_0$. On the other hand, if $u = e^{-\lambda\tau/2} R$ for some fixed $R > 0$, then

$$\left| \frac{e^{-2\lambda\tau} Z_2(u)}{e^{-\lambda\tau} Z_1(u)} \right| \lesssim c \left(R^{-2} + e^{-\lambda\tau} O(\log R + \tau) \right),$$

which is small for all sufficiently large τ if for a given c we choose R to be large.

Let us label the region where $ue^{\lambda\tau/2} = O(1)$ as the *interior region*. The complement of the interior region is labelled as the *exterior region*. Then in the exterior region, the formal solution is

$$z_{\text{form, ext}} = cu^{-2} (1 + u^2)^{1-\lambda} e^{-\lambda\tau}.$$

3.2. Formal solution in the interior region. In the interior region where $ue^{\lambda\tau/2} = O(1)$, we introduce a new variable

$$r := ue^{\lambda\tau/2}.$$

Then in the (r, τ) -coordinates, since

$$\begin{aligned} \partial_\tau|_r z &= \partial_\tau|_u z - \frac{\lambda}{2} u z_u = \partial_\tau|_u z - \frac{\lambda}{2} r z_r, \\ \mathcal{E}_u[z] &= e^{\lambda\tau} \mathcal{E}_r[z], \end{aligned}$$

equation (3.1) becomes

$$(3.6) \quad e^{-\lambda\tau} \left\{ \partial_\tau|_r z + \frac{\lambda-1}{2} r z_r \right\} - \frac{1}{2(n-1)} \mathcal{E}_r[z] = 0.$$

Suppose, for the sake of the formal argument, that the term involving $e^{-\lambda\tau}$ is negligible for sufficiently large τ , then this equation is approximated by the equation

$$\mathcal{E}_r[\tilde{z}] = 0,$$

whose solution, subject to the boundary $\tilde{z}(0) = 1$ and the asymptotic condition $\lim_{r \nearrow \infty} \tilde{z}(r) = 0$, is a Bryant soliton profile function

$$\tilde{z}(r) = \mathfrak{B}(Ar),$$

where A does not depend on r . The complete smooth metric given by

$$g = \mathfrak{B}^{-1}(Ar) dr^2 + r^2 g_{\text{sph}}$$

is a scaled version of the Bryant soliton [7].

The function $\mathfrak{B}(r)$ is smooth and strictly monotonically decreasing for all $r > 0$ with the following asymptotics

$$(3.7) \quad \mathfrak{B}(r) = \begin{cases} 1 - b_2 r^2 + b_3 r^4 + b_3 r^6 + \dots, & r \searrow 0, \\ c_1 r^{-2} + c_2 r^{-4} + c_3 r^{-6} + \dots, & r \nearrow \infty, \end{cases}$$

where b_k 's and c_k 's are constants; in particular, $b_2 > 0$. In this paper, we normalize $\mathfrak{B}(r)$ by setting $c_1 = 1$. In the interior region, our formal solution is

$$z_{\text{form, int}} = \mathfrak{B}(Ar).$$

Remark 3.1. If $\lambda = 1$, then $r = ue^{\tau/2} = \psi/\sqrt{2(n-1)}$. In this case, the Bryant soliton $\mathfrak{B}(Ar) = \mathfrak{B}\left(A\psi/\sqrt{2(n-1)}\right)$ solves equation (2.5) and gives a trivial example of Ricci flow with Type-IIb singularity with the highest curvature blowing up at the rate $O(t^{\lambda-1}) = O(\lambda^0) = O(1)$. In comparison, we construct different solutions in Theorem 1.1 for the case $\lambda = 1$.

3.3. Matching condition. We now match the formal solutions at the interface of the interior region and the exterior region. If we pick $r = R \gg 1$, then in the interior region, using the asymptotics of $z_{\text{form, int}}$ as $r \nearrow \infty$, we have

$$(3.8) \quad z_{\text{form, int}} = \mathfrak{B}(AR) \approx A^{-2}R^{-2};$$

in the exterior region, using the asymptotics of $z_{\text{form, ext}}$ as $u \searrow 0$ and that $u = Re^{-\lambda\tau/2}$, we have

$$(3.9) \quad z_{\text{form, ext}} = cu^{-2}(1+u^2)^{1-\lambda}e^{-\lambda\tau} \approx cR^{-2}.$$

Equating (3.8) and (3.9), we obtain the matching condition for the formal solution

$$(3.10) \quad A^{-2} = c.$$

The condition 3.10 says that given A and $R > 0$, we can always find such c ; equivalently, fixing c and R , then A is determined.

3.4. Features of the formal solution. Our formal solutions defined in the interior region and the exterior region are valid for all dimensions $n+1 \geq 3$ and give rise to Riemannian metrics on \mathbb{R}^{n+1} as defined in (2.4). In fact, these Riemannian metrics are complete, as will be proven in Lemma 6.1. Since $z_{\text{form, ext}} = e^{-\lambda\tau}cu^{-2}(1+u^2)^{1-\lambda}$, we have at any $\tau < \infty$, as $u \nearrow \infty$, i.e., $\psi = \sqrt{2(n-1)}ue^{\tau/2} \nearrow \infty$, that

$$(3.11) \quad z = \psi_s^2 \sim e^{-\lambda\tau}cu^{-2\lambda} \sim c\psi^{-2\lambda},$$

$$(3.12) \quad K = -\frac{z\psi}{2\psi} \sim -\frac{z_u}{2u}e^{-\tau} \sim (tu^2)^{-(1+\lambda)} \sim \psi^{-2(1+\lambda)},$$

$$(3.13) \quad L = \frac{1-z}{\psi^2} = \frac{1-z}{u^2}e^{-\tau} \sim (tu^2)^{-1} \sim \psi^{-2}.$$

Remark 3.2. Conditions (3.12) and (3.13) imply that the metrics given by the barriers (cf. Section 5), and hence the Ricci flow solutions described in Theorem 1.1 (cf. Section 6), are in fact asymptotically flat in the sense that $|\text{Rm}| \rightarrow 0$ as one approaches spatial infinity. Recall that an asymptotically conical metric on \mathbb{R}^{n+1} is given by $ds^2 + \alpha^2 s^2 g_{\text{sph}}$, where $\alpha \in (0, 1]$ with

the case $\alpha = 1$ being the Euclidean metric. For an asymptotically conical metric, $z = \psi_s^2 = \alpha^2$. As will be shown in Section 6, the solutions in Theorem 1.1 do not satisfy $z \equiv 1$ and therefore are non-Euclidean. Also, condition (3.11) implies that the metric $ds^2 + \psi^2 g_{\text{sph}}$ we construct in this paper are *not* asymptotically Euclidean in the sense considered in [12] or [25].

As we move towards the origin \mathcal{O} , $z(u) \nearrow 1$ and we enter the interior region where the formal solution z_{form} is a Bryant soliton profile function. At \mathcal{O} , we have $K(\mathcal{O}, t) = L(\mathcal{O}, t)$ for all $t \geq t_0$, and the norm of the curvature tensor achieves its maximum value

$$|\text{Rm}(\mathcal{O}, t)| = \sqrt{n(n+1)}L(\mathcal{O}, t) = \frac{\sqrt{n(n+1)}}{2(n-1)} \lim_{r \searrow 0} \frac{1-z}{r^2} e^{(\lambda-1)\tau} = t^{\lambda-1}C,$$

where C is a positive constant depending on n, A, b_2 ; to be precise, $C = \frac{\sqrt{n(n+1)}A^2b_2}{2(n-1)}$. Therefore, the formal solution has a Type-IIIb singularity if $\lambda > 0$. In particular, the curvature of a Ricci flow solution that asymptotically approaches this formal solution necessarily blows up at the same rate.

4. SUBSOLUTIONS AND SUPERSOLUTIONS

Given a parabolic differential operator $\mathcal{P}[v] = \partial_\tau v - \mathcal{D}[v]$ where $\mathcal{D}[\cdot]$ is some second-order elliptic operator, a function v^+ is a *subsolution* of the PDE $\mathcal{P}[v] = 0$ if $\mathcal{D}[v^+] \leq 0$ whereas a function v^- is a *supersolution* if $\mathcal{D}[v^-] \geq 0$. If there exist subsolution v^- and supersolution v^+ and in addition, $v^- \leq v^+$, then we call v^- a *lower barrier* and v^+ an *upper barrier*.

Suppose the equation $\mathcal{P}[v] = 0$ admits a solution, then the existence of barriers $v^- \leq v^+$ implies that there exists a solution v with $v^- \leq v \leq v^+$. This is the general idea of our argument which will be justified rigorously in this section and next. In this section, we construct subsolutions and supersolutions for equation (3.1) in the interior and the exterior regions. In the next section, we patch them to obtain global upper and lower barriers.

4.1. Interior region. Let us define

$$(4.1) \quad \mathcal{T}_r[z] := e^{-\lambda\tau} \left\{ \partial_\tau|_r z + \frac{\lambda-1}{2} r z_r \right\} - \frac{1}{2(n-1)} \mathcal{E}_r[z].$$

Then according to (3.6), $z(r, \tau)$ satisfies the equation $\mathcal{T}_r[z] = 0$ in the interior region. The subsolution and supersolution for this equation are given in the next lemma.

Lemma 4.1. *For an integer $n \geq 2$, a real number $\lambda > 0$ and a constant $A > 0$, there exist a sufficiently large $\tau_1 < \infty$, a constant $B_1 > 0$, a bounded function $\beta(r, \tau) : (0, \infty) \times [\tau_1, \infty) \rightarrow \mathbb{R}$ and constants a^\pm , all depending only on A , such that the functions*

$$(4.2) \quad z_{\text{int}}^\pm(r, \tau) := \mathfrak{B} \left(A \left(1 + a^\pm e^{-\lambda\tau/2} \right) r \right) \pm e^{-\lambda\tau} \beta(r, \tau)$$

are supersolution (+) and subsolution (-), respectively, of $\mathcal{J}_r[z] = 0$ in the region $\Omega_{int} := \{0 \leq r \leq B_1 e^{\lambda\tau/2}\}$ for all $\tau \geq \tau_1$.

Proof. Let us denote $\mathbf{B}^\pm(r, \tau) := \mathfrak{B}(A(1 + a^\pm e^{-\lambda\tau/2})r)$. Then

$$\partial_\tau|_r \mathbf{B}^\pm = r \mathbf{B}_r \frac{-\lambda a^\pm e^{-\lambda\tau/2}}{2(1 + a^\pm e^{-\lambda\tau/2})}.$$

In order for $z_{int}^+ = \mathbf{B}^+(r, \tau) + e^{-\lambda\tau} \beta(r, \tau)$ to be a supersolution, we need to show $\mathcal{J}_r[z_{int}^+] \geq 0$. In below, for notational clarity, we drop the superscript “+”.

Since $\mathbf{B}(r, \tau)$ solves $\mathcal{E}_r[z] = 0$, we obtain

$$\begin{aligned} \mathcal{J}_r[z_{int}^+] &= e^{-\lambda\tau} \left\{ -\frac{\mathcal{L}_r[\beta] + 2\hat{\mathcal{Q}}_r[\mathbf{B}, \beta]}{2(n-1)} + \frac{\lambda-1}{2} r \mathbf{B}_r \right\} \\ &\quad + e^{-3\lambda\tau/2} r \mathbf{B}_r \frac{-\lambda a}{2(1 + a e^{-\lambda\tau/2})} \\ &\quad + e^{-2\lambda\tau} \left\{ -\lambda\beta + \partial_\tau|_r \beta + \frac{\lambda-1}{2} r \beta_r - \frac{\mathcal{Q}_r[\beta]}{2(n-1)} \right\}. \end{aligned}$$

Set $\hat{A} := 1 + \frac{\lambda-1}{2} = \frac{\lambda+1}{2} > 0$, we define $\beta(r, \tau)$ to be a solution of the equation

$$(4.3) \quad \mathcal{L}_r[\beta] + 2\hat{\mathcal{Q}}_r[\mathbf{B}, \beta] = 2(n-1)\hat{A}r\mathbf{B}_r.$$

Using the definitions of \mathcal{L}_r in (2.6) and $\hat{\mathcal{Q}}_r$ in (2.7) respectively, equation (4.3) becomes

$$(4.4) \quad \mathbf{B}\beta_{rr} + \left\{ \frac{n-1}{r} - \mathbf{B}_r - \frac{\mathbf{B}}{r} \right\} \beta_r + \left\{ \mathbf{B}_{rr} - \frac{\mathbf{B}_r}{r} + 2(n-1)\frac{1-2\mathbf{B}}{r^2} \right\} \beta = 2(n-1)\hat{A}r\mathbf{B}_r.$$

Using the asymptotic expansions of $\mathbf{B}(r, \tau)$ near $r = 0$ and $r = \infty$ given in (3.7), we have the following. Near $r = 0$, equation (4.4) is approximated by

$$\beta_{rr} + \frac{n-2}{r}\beta_r - \frac{2(n-1)}{r^2}\beta = -C_1 r^2 \left(1 + O\left(ae^{-\lambda\tau/2}\right)\right),$$

where $C_1 = 2(n-1)(\gamma+1)b_2A^2$. Near $r = \infty$, equation (4.4) is a perturbation of the equation

$$\frac{1 + O\left(ae^{-\lambda\tau/2}\right)}{(Ar)^2} \beta_{rr} + \frac{n-1}{r}\beta_r + \frac{2(n-1)}{r^2}\beta = -\frac{4(n-1)\hat{A}}{(Ar)^2} \left(1 + O\left(ae^{-\lambda\tau/2}\right)\right).$$

So there exists a solution β to equation (4.3) with the following asymptotics

$$(4.5) \quad \beta(r, \tau) = \begin{cases} r^2 + O\left(r^4(1 + ae^{-\lambda\tau/2})\right), & r \searrow 0, \\ \left(-2\hat{A}/A^2 + o(1)\right) \left(1 + O\left(ae^{-\lambda\tau/2}\right)\right), & r \nearrow \infty. \end{cases}$$

Also, the asymptotic expansions

$$-r\mathbf{B}_r = \begin{cases} (C_7 r^2 + o(r^2)) (1 + O(ae^{-\lambda\tau/2})), & r \searrow 0, \\ (C_8 r^{-2} + o(r^{-2})) (1 + O(ae^{-\lambda\tau/2})), & r \nearrow \infty, \end{cases}$$

imply that

$$-r\mathbf{B}_r \geq C_9 \min\{r^2, r^{-2}\}.$$

Then in view of (4.5), we have for $0 < r \leq 1$,

$$\left| -\lambda\beta + \partial_\tau|_r \beta + \frac{\lambda-1}{2} r\beta_r - \frac{\mathcal{Q}_r[\beta]}{2(n-1)} \right| \leq C_{10} r^2,$$

and hence

$$\begin{aligned} \mathcal{J}_r [z_{\text{int}}^+] &\geq -e^{-\lambda\tau} r\mathbf{B}_r - e^{-3\lambda\tau/2} C_7 r^2 - e^{-2\lambda\tau} C_{10} r^2 \\ &\geq e^{-\lambda\tau} r^2 \left(C_9 - e^{-\lambda\tau/2} C_7 - e^{-\lambda\tau} C_{10} \right) \\ &> 0 \end{aligned}$$

for all $\tau \geq \tau_1$ with τ_1 sufficiently large. And for $r \geq 1$,

$$\left| -\lambda\beta + \partial_\tau|_r \beta + \frac{\lambda-1}{2} r\beta_r - \frac{\mathcal{Q}_r[\beta]}{2(n-1)} \right| \leq C_{11},$$

so then

$$\begin{aligned} \mathcal{J}_r [z_{\text{int}}^+] &\geq -e^{-\lambda\tau} r\mathbf{B}_r - e^{-3\lambda\tau/2} C_8 r^{-2} - e^{-2\lambda\tau} C_{11} \\ &\geq e^{-\lambda\tau} \left(C_9 r^{-2} - e^{-\lambda\tau/2} C_8 r^{-2} - e^{-\lambda\tau} C_{11} \right) \\ &> 0 \end{aligned}$$

provided that $r < B_1 e^{\lambda\tau/2}$ with constant $B_1 := \sqrt{C_9/(2C_{11})}$.

Therefore, z_{int}^+ is indeed a supersolution. That z_{int}^- is a subsolution is proved similarly. So the lemma follows. \square

Remark 4.2. For all $\tau \geq \tau_1$, $z_{\text{int}}^- < \mathfrak{B}(Ar) < z_{\text{int}}^+$ in Ω_{int} .

4.2. Exterior region. Let us define

$$(4.6) \quad \mathcal{F}_u[z] := \partial_\tau|_u z - \frac{1}{2(n-1)} \mathcal{E}_u[z] - \frac{1}{2} u z_u.$$

By definition, $\mathcal{E}_u[z] = \mathcal{L}_u[z] + \mathcal{Q}_u[z]$, so

$$\mathcal{F}_u[z] = \partial_\tau|_u z - \frac{1}{2} (u^{-1} + u) - u^{-2} z - \frac{\mathcal{Q}_u[z]}{2(n-1)}.$$

Then in this region, $z(u, \tau)$ satisfies the equation $\mathcal{F}_u[z] = 0$, cf. (3.1). The next lemma takes care of the subsolution and supersolution for this equation.

From now on, let us define $Z(u) := u^{-2} (1 + u^2)^{1-\lambda}$. We note that $Z(u) > 0$ for all $u \in (0, \infty)$.

Lemma 4.3. *For an integer $n \geq 2$, a real number $\lambda > 0$ and constants $c^\pm > 0$, there exist function $\zeta : (0, \infty) \rightarrow \mathbb{R}$, constants $B_2^\pm > 0$, a sufficiently large $\tau_2 < \infty$, and a constants b_*^\pm depending only on c^\pm , respectively, such that for any $b \geq b_*$, the functions*

$$(4.7) \quad z_{\text{ext}}^\pm(u, \tau) := c^\pm e^{-\lambda\tau} Z(u) \pm b^\pm e^{-2\lambda\tau} \zeta(u)$$

are supersolution (+) and subsolution (-), respectively, of $\mathcal{F}_u[z] = 0$ in the region $\Omega_{\text{ext}}^\pm := \left\{ B_2 \sqrt{\frac{b^\pm}{c^\pm}} e^{-\lambda\tau/2} \leq u < \infty \right\}$ and for all $\tau \geq \tau_2$.

Proof. We first prove the lemma for z_{ext}^+ . To simplify notation, we omit the superscript “+” in the argument below.

Since $Z(u)$ is a solution of the ODE (3.3), we have

$$\begin{aligned} e^{2\lambda\tau} \mathcal{F}_u [z_{\text{ext}}^+] &= b \left\{ -\frac{1}{2} (u^{-1} + u) \zeta' - (u^{-2} + 2\lambda) \zeta \right\} - \frac{c^2}{2(n-1)} \mathcal{Q}_u[Z] \\ &\quad - \frac{bc}{n-1} e^{-\lambda\tau} \hat{\mathcal{Q}}_u[Z, \zeta] - \frac{b^2}{2(n-1)} e^{-2\lambda\tau} \mathcal{Q}_u[\zeta]. \end{aligned}$$

Since

$$\mathcal{Q}_u [Z] = 2u^{-6} (1 + u^2)^{-2\lambda} f(u),$$

where

$$f(u) = 4 - n(1 + u^2)^2 + u^4(1 + \lambda)^2 + 2u^2(3 + \lambda),$$

we have for $u \in (0, \infty)$,

$$(4.8) \quad |f(u)| \leq C_1(1 + u^2)^2$$

for some constant C_1 depending only on n and λ .

Let $\zeta : (0, \infty) \rightarrow \mathbb{R}$ be a solution of the ODE

$$(4.9) \quad -\frac{1}{2} (u^{-1} + u) \zeta' - (u^{-2} + 2\lambda) \zeta = u^{-6} (1 + u^2)^{2-2\lambda}.$$

Then we solve this ODE to obtain

$$\zeta(u) := u^{-4} (1 + u^2)^{1-2\lambda} (1 + C_2 u^2)$$

for an arbitrary constant C_2 . Let us choose $C_2 = 1$, so

$$(4.10) \quad \zeta(u) := u^{-4} (1 + u^2)^{2-2\lambda}.$$

In particular, $\zeta(u) > 0$ for all $u \in (0, \infty)$.

From (4.10), the asymptotics of ζ are

$$\zeta(u) = \begin{cases} u^{-4} + O(u^{-2}), & u \searrow 0, \\ C_4 u^{-4\lambda} + O(u^{-2-4\lambda}), & u \nearrow \infty. \end{cases}$$

So the following estimates hold. For $B_2 e^{-\lambda\tau/2} \leq u < 1$,

$$\left| \hat{\mathcal{Q}}_u [Z, \zeta] \right| \leq C_5 u^{-8}, \quad \left| \mathcal{Q}_u [\zeta] \right| \leq C_6 u^{-10}$$

For $1 \leq u < \infty$,

$$\left| \hat{\mathcal{Q}}_u [Z, \zeta] \right| \leq C_7 u^{-2-6\lambda}, \quad |\mathcal{Q}_u [\zeta]| \leq C_8 u^{-2-8\lambda}.$$

Using the definition of ζ and estimate (4.8), we have

$$\begin{aligned} e^{2\lambda\tau} \mathcal{F}_u [z_{\text{ext}}^+] &= \left(b - \frac{C_1 c^2}{n-1} \right) u^{-6} (1+u^2)^{2-2\lambda} \\ &\quad - \frac{bc}{n-1} e^{-\lambda\tau} \hat{\mathcal{Q}}_u [Z, \zeta] - \frac{b^2}{2(n-1)} e^{-2\lambda\tau} \mathcal{Q}_u [\zeta] \\ &\geq \frac{b - C_1 c^2}{n-1} u^{-6} (1+u^2)^{2-2\lambda} \\ &\quad - \frac{bc}{n-1} e^{-\lambda\tau} \left| \hat{\mathcal{Q}}_u [Z, \zeta] \right| - \frac{b^2}{2(n-1)} e^{-2\lambda\tau} |\mathcal{Q}_u [\zeta]|. \end{aligned}$$

We choose $b_* = c^2 (1 + C_1/(n-1))$. Then for any $b \geq b_*$, we have the following. Then for $0 < u \leq 1$, there exists a constant B_2 such that

$$\begin{aligned} e^{2\lambda\tau} \mathcal{F}_u [z_{\text{ext}}^+] &\geq C_1 u^{-6} \left(c^2 - C_5 b c u^{-2} e^{-\lambda\tau} - C_6 b^2 u^{-4} e^{-2\lambda\tau} \right) \\ &\geq C_1 u^{-6} \left(c^2 - C_5 c^2 B_2^{-2} - C_6 c^2 B_2^{-2} \right) \\ &\geq \frac{C_1}{2} u^{-6} c^2 \\ &> 0 \end{aligned}$$

provided that $u^2 e^{\lambda\tau} \geq B_2^2 b/c$, or equivalently,

$$B_2 \sqrt{\frac{b}{c}} e^{-\lambda\tau/2} \leq u \leq 1.$$

For $1 \leq u < \infty$, since

$$\left| u^{-6} (1+u^2)^{2-2\lambda} \right| \leq C_1 u^{-2-4\lambda},$$

we have

$$\begin{aligned} e^{2\lambda\tau} \mathcal{F}_u [z_{\text{ext}}^+] &\geq C_1 u^{-2-4\lambda} \left(c^2 - C_5 c u^{-2\lambda} e^{-\lambda\tau} - C_6 c^2 u^{-4\lambda} e^{-2\lambda\tau} \right) \\ &\geq C_1 u^{-2-4\lambda} \left(c^2 - C_5 b c e^{-\lambda\tau} - C_6 b^2 e^{-2\lambda\tau} \right) \\ &> 0 \end{aligned}$$

for all $\tau \geq \tau_2$ with τ_2 sufficiently large.

Therefore, z_{ext}^+ is indeed a supersolution. By a similar argument, z_{ext}^- is a subsolution. So the lemma is proven. \square

Remark 4.4. If we let $c^- \leq c \leq c^+$, then for all $\tau \geq \tau_2$, $z_{\text{ext}}^- < e^{-\lambda\tau} cZ(u)$ in Ω_{ext}^- and $e^{-\lambda\tau} cZ(u) < z_{\text{ext}}^+$ in Ω_{ext}^+ .

5. UPPER AND LOWER BARRIERS

According to Lemmata 4.1 and 4.3, the interior region Ω_{int} and the exterior region Ω_{ext}^\pm overlap for sufficiently large τ . Our goal in this section is to show that the regional supersolutions z_{int}^+ and z_{ext}^+ together with z_{int}^- and z_{ext}^- can be patched together to provide an upper and lower barriers, respectively, for Ricci flow equation (3.1).

In the next two lemmata, we prove in each region the subsolution and supersolution are ordered.

Lemma 5.1. *Let $\beta(r, \tau)$ be defined as in Lemma 4.1. For $a^- > a^+$, there exists $\tau_3 \geq \tau_1$ such that*

$$z_{\text{int}}^\pm := \mathfrak{B} \left(A \left(1 + a^\pm e^{-\lambda\tau/2} \right) r \right) \pm e^{-\lambda\tau} \beta(r, \tau)$$

satisfy $\lambda_{\text{int}}^- < \lambda_{\text{int}}^+$ in $\{0 \leq r \leq B_1 e^{\lambda\tau/2}\}$ for all $\tau \geq \tau_3$.

Proof. Using the asymptotic expansions of \mathfrak{B} (3.7) and β (4.5), we have the following for all sufficiently large $\tau \geq \tau_1$. Near $r = 0$,

$$\begin{aligned} z_{\text{int}}^+ - z_{\text{int}}^- &= e^{-\lambda\tau/2} (2b_2 A^2 (a^- - a^+) r^2 + O(r^{-4})) + e^{-\lambda\tau} (-2\hat{A}A^{-2} + o(1)) \\ &> 0. \end{aligned}$$

Near $r = \infty$, with $\hat{A} = (\gamma + 1)/2$,

$$\begin{aligned} z_{\text{int}}^+ - z_{\text{int}}^- &= e^{-\lambda\tau/2} (2A^{-2} (a^- - a^+) r^{-2} + O(r^4)) + e^{-\lambda\tau} (r^2 + o(r^2)) \\ &> 0. \end{aligned}$$

On any bounded interval $c < r < C$, it is straightforward to check that $\lambda_{\text{int}}^- < \lambda_{\text{int}}^+$. So the lemma is proved. \square

Lemma 5.2. *Let $R_2 := \max \left\{ B_2^+ \sqrt{\frac{b^+}{c^+}}, B_2^- \sqrt{\frac{b^-}{c^-}} \right\}$. If $c^+ \geq c^-$, then*

$$z_{\text{ext}}^\pm := c^\pm e^{-\lambda\tau} Z(u) \pm b^\pm e^{-2\lambda\tau} \zeta(u)$$

satisfy $\lambda_{\text{ext}}^- < \lambda_{\text{ext}}^+$ in $\{R_2 e^{-\lambda\tau/2} < u < \infty\}$ for all $\tau \geq \tau_2$.

Proof. Using the definitions of Z and ζ , and choosing $C_2 \geq 0$, and recall that $c^+ \geq c^-$ implies $b^+ > b^-$, we have

$$\begin{aligned} e^{\lambda\tau} (z_{\text{ext}}^+ - z_{\text{ext}}^-) &= (c^+ - c^-) Z(u) + e^{-\lambda\tau} (b^+ + b^-) \zeta(u) \\ &= (c^+ - c^-) \frac{(1 + u^2)^{1+\lambda}}{u^2} + e^{-\lambda\tau} \frac{(b^+ + b^-) (1 + (n-1)C_2 u^2)}{(n-1)u^4 (1 + u^2)^{2\lambda-1}} \\ &> 0 \end{aligned}$$

for all $u \in (0, \infty)$ for all $\tau \geq \tau_2$. So the lemma is proved. \square

To patch the supersolution in the interior region with that in the exterior region, we state and prove a patching lemma for z_{int}^+ and z_{ext}^+ . We omit the patching lemma for z_{int}^- and z_{ext}^- , since its statement and proof are analogous. To shorten the notation, we write a^+, b^+, c^+ as a, b, c .

Lemma 5.3. *Let $R_D := D\sqrt{b/c}$ where $D > 0$ is arbitrary. Suppose A , a and c satisfy the following inequality*

$$(5.1) \quad \left(1 + \frac{3}{8}D^{-2}\right)c < A^{-2} < \left(1 + \frac{1}{2}D^{-2}\right)c.$$

Then there exists $\tau_5 \geq \max\{\tau_3, \tau_4\}$ sufficiently large such that

$$(z_{\text{int}}^+ - z_{\text{ext}}^+)(R_D) < 0, \quad (z_{\text{int}}^+ - z_{\text{ext}}^+)(2R_D) > 0$$

for all $\tau \geq \tau_4$.

Proof. At the interface of the interior and exterior regions, we have the following for $\tau \geq \tau_5$. In the interior region, we have as $r \nearrow \infty$ that

$$\begin{aligned} z_{\text{int}}^+ &= \mathbf{B}(r, \tau) + e^{-\lambda\tau}\beta(r, \tau) \\ &= (A^{-2}r^{-2} + c_2A^{-4}r^{-4} + O(r^{-6}))\left(1 + O(ae^{-\lambda\tau/2})\right) + O(e^{-\lambda\tau}). \end{aligned}$$

In the exterior region, we have as $u = re^{-\lambda\tau/2} \searrow 0$ that

$$\begin{aligned} z_{\text{ext}}^+ &= e^{-\lambda\tau}(cu^{-2} + O(1)) + e^{-2\lambda\tau}(bu^{-4} + O(u^{-2})) \\ &= cr^{-2} + br^{-4} + O(e^{-\lambda\tau}r^{-2}) \end{aligned}$$

So on bounded r -interval, we have

$$r^2(z_{\text{int}}^+ - z_{\text{ext}}^+) = (A^{-2} - c) + (c_2A^{-4} - b + O(r^{-2}))r^{-2} + O(e^{-\lambda\tau/2}).$$

Let us choose a constant \hat{C} so large that for

$$b \geq \hat{C}A^{-4} \quad \text{and} \quad b \geq \hat{C}\sqrt{c},$$

we have

$$\left|\frac{c_2c}{bA^4} + O\left(\frac{c^2}{b^2}\right)\right| \leq \frac{c}{2}.$$

Then at $r = R_D$,

$$\begin{aligned} R_D^2(z_{\text{int}}^+ - z_{\text{ext}}^+) &= (A^{-2} - c) + \left[\frac{c_2c}{bA^4} + O\left(\frac{c^2}{b^2}\right) - c\right]D^{-2} + O(\tau e^{-\lambda\tau}) \\ &\leq A^{-2} - \left(1 + \frac{1}{2}D^{-2}\right)c + O(e^{-\lambda\tau/2}), \end{aligned}$$

and at $r = 2R_D$,

$$\begin{aligned} 4R_D^2(z_{\text{int}}^+ - z_{\text{ext}}^+) &= (A^{-2} - c) + \left[\frac{c_2c}{bA^4} + O\left(\frac{c^2}{b^2}\right) - c\right]\frac{D^{-2}}{4} + O(\tau e^{-\lambda\tau}) \\ &\geq A^{-2} - \left(1 + \frac{3}{8}D^{-2}\right)c + O(e^{-\lambda\tau/2}). \end{aligned}$$

Now choose A and c according to (5.1), then the lemma follows for $\tau \geq \tau_5$. \square

For fixed $\lambda \neq 1$ and constants A, b^\pm, c^\pm chosen so far, we define the upper barrier z^+ for equation (3.1) by

$$(5.2) \quad z^+ := \begin{cases} z_{\text{int}}^+, & \text{if } 0 < u \leq R_D e^{-\lambda\tau/2}, \\ \inf \{z_{\text{int}}^+, z_{\text{ext}}^+\}, & \text{if } R_D e^{-\lambda\tau/2} \leq u \leq 2R_D e^{-\lambda\tau/2}, \\ z_{\text{ext}}^+, & \text{if } 2e^{-\lambda\tau/2} R_D \leq u < \infty. \end{cases}$$

The lower barrier $z^- = z^-(u, \tau)$ for equation (3.1) is defined analogously using z_{int}^- and z_{ext}^- ; in particular, $z^- := \sup\{z_{\text{int}}^-, z_{\text{ext}}^-\}$ for $R_D e^{-\lambda\tau/2} \leq u \leq 2R_D e^{-\lambda\tau/2}$. By remarks 4.2 and 4.4, we see that z^+ stays strictly above the formal solution and z^- strictly below the formal solution.

Lemmata 5.1–5.3 imply the following proposition.

Proposition 5.4. *There exist a sufficiently large $\tau_0 < \infty$ and positive continuous, piecewise smooth functions $z^\pm = z^\pm(u, \tau)$ defined for $0 < u < \infty$ and $\tau \geq \tau_0$ such that the following hold.*

- (B1) z^\pm are upper (+) and lower (–) barriers to equation (3.1), respectively.
- (B2) $z^- < z^+$; near $u = 0$, $z^\pm = z_{\text{int}}^\pm$; as $u \nearrow \infty$, $z^\pm = z_{\text{ext}}^\pm$.
- (B3) At any $\tau \in [\tau_0, \infty)$, we have

$$\lim_{u \searrow 0} z^- = \lim_{u \searrow 0} z^+ = 1, \quad \lim_{u \nearrow \infty} z^- = \lim_{u \nearrow \infty} z^+ = 0.$$

Remark 5.5. By construction, where z^+ (or z^-) is not smooth, the corner is concave (or convex).

We end this section with a comparison principle for the equation (3.1).

Proposition 5.6. *Let $\bar{\tau} \in [\tau_0, \infty)$ be arbitrary. Let z^\pm be two non-negative sub-(–) and super-(+) solutions of equation (3.1) respectively. Moreover, assume*

- (C1) $z^-(u, \tau_0) < z^+(u, \tau_0)$ for $0 < u < \infty$;
- (C2) $z^-(0, \tau) \leq z^+(0, \tau)$, and $\lim_{u \nearrow \infty} (z^-(u, \tau) - z^+(u, \tau)) \leq 0$ for all $\tau \in [\tau_0, \bar{\tau}]$.

Then $z^-(u, \tau) \leq z^+(u, \tau)$ in $[0, \infty) \times [\tau_0, \bar{\tau}]$.

Remark 5.7. In this proposition, we assume z^\pm are smooth. The result also holds for the continuous, piecewise smooth barriers z^\pm constructed earlier. When applying the comparison principle, we will only evaluate z^\pm at “points of first contact with a given smooth function” which are necessarily smooth points of z^\pm for each $\tau \geq \tau_0$.

Proof of Proposition 5.6. By (C1) and (C2), for any given $\varepsilon > 0$, there exists $R = R(\Delta)$ such that $z^+ > z^-$ on $(0, R] \times [\tau, \bar{\tau}]$ and $(z^+ - z^-)(R) > \varepsilon$.

Define

$$w := e^{-\mu\tau} (z^+ - z^-) + \varepsilon,$$

where $\mu > 0$ is to be chosen. Then $w > 0$ on the parabolic boundary of the evolution by assumptions (C1) and (C2). We claim that $w > 0$ in $(0, R) \times [\tau_0, \bar{\tau}]$. Suppose the contrary, then there must be an interior point u_* and a first time τ_* such that $w(u_*, \tau_*) = 0$ and $w_\tau(u_*, \tau_*) \leq 0$. Moreover, at (u_*, τ_*) , we have

$$z^+ = z^- - \epsilon e^{-\mu\tau}, \quad z_u^+ = z_u^-, \quad z_{uu}^+ \geq z_{uu}^-.$$

Then at (u_*, τ_*) ,

$$\begin{aligned} 0 &\geq e^{\mu\tau_*} \partial|_\tau w_\tau \\ &= (z_\tau^+ - z_\tau^-) - \mu(z^+ - z^-) \\ &\geq (z^+ - z^-)(u^{-2} - \mu) + \frac{\mathcal{Q}_u[z^+] - \mathcal{Q}_u[z^-]}{2(n-1)} \\ &= (z^- - z^+) \left\{ \mu + \frac{(z_u^+/u) - z_{uu}^+}{2(n-1)} + \frac{z^+ + z^- - 1}{u^2} \right\} + z^- (z_{uu}^+ - z_{uu}^-) \\ &\geq \epsilon e^{-\mu\tau_*} \left\{ \mu - \frac{\mathcal{Q}_u[z^+] - \mathcal{Q}_u[z^-]}{2(n-1)} \Big|_{(u_*, \tau_*)} - \frac{1}{u_*^2} \right\} \\ &= \epsilon e^{-\mu\tau_*} \{ \mu - (\text{bounded term independent of } \mu) \} \end{aligned}$$

Since $\epsilon > 0$ is fixed, we choose μ sufficiently large, then at (u_*, τ_*) we have

$$0 \geq \partial|_\tau w > 0,$$

which is a contradiction. Hence, the claim is true. In the proof of the claim, μ may depend on ζ^+ , ζ^- and $\bar{\tau}$, but not on $\epsilon > 0$. Therefore, letting $\epsilon \rightarrow 0$, the proposition follows. \square

6. PROOF OF THEOREM 1.1

For any solution z of equation (3.1) we have the following.

Lemma 6.1. *Suppose $0 < z \leq z^+$. If $\lambda > 0$, then z determines a complete rotationally symmetric metric $g := z^{-1}d\psi^2 + \psi^2 g_{sph}$ on \mathbb{R}^N .*

Proof. By definition g is rotationally symmetric. To see that g is a complete metric, it suffices to show that any radial geodesic γ starting from the origin has infinite length in the s -coordinate. The length of γ in s -coordinate is a function of u and τ given by

$$s(u, \tau) = e^{-\tau/2} \sigma(u) = e^{-\tau/2} \int_0^u \frac{d\sigma}{d\hat{u}} d\hat{u}.$$

Since $z = \psi_s^2 = 2(n-1)u_\sigma^2$, and $0 < z \leq z^+$ by hypothesis, we have

$$\frac{\sigma(u)}{\sqrt{2(n-1)}} \geq \int_{u_0}^u \frac{1}{\sqrt{z}} d\hat{u} \geq \int_{u_0}^u \frac{1}{\sqrt{z^+}} d\hat{u}.$$

Recall that

$$z_{\text{ext}}^+ = e^{-\lambda\tau} c u^{-2} (1 + u^2)^{1-\lambda} + e^{-2\lambda\tau} b u^{-4} (1 + u^2)^{2-2\lambda}$$

So for u_0 and τ_0 sufficiently large, $z^+ = z_{\text{ext}}^+$ in $[u_0, 1) \times [\tau_0, \infty)$ with

$$z_{\text{ext}}^+ \lesssim e^{-\lambda\tau} u^{-2\lambda}.$$

It follows that

$$e^{\tau/2} s(u, \tau) \gtrsim \int_{u_0}^u \frac{1}{\sqrt{z^+}} d\hat{u} = \int_{u_0}^u \frac{1}{\sqrt{z_{\text{ext}}^+}} d\hat{u} \gtrsim e^{\lambda\tau/2} \int_{u_0}^u \hat{u}^\lambda d\hat{u} = u^{1+\lambda} - u_0^{1+\lambda}.$$

So for each $\tau \geq \tau_0$, $\lim_{u \nearrow \infty} s(u, \tau) = \infty$, whence the lemma follows. \square

Since $z = \psi_s^2$, where s is the arclength from the origin, and we are working with complete metrics on \mathbb{R}^{n+1} , we have $\psi_s > 0$. Also, our formal solution and barriers all satisfy $\lim_{s \rightarrow 0^+} \psi_s = 1$ and $\lim_{s \nearrow \infty} \psi_s = 0$. So we have the following Lemma which is a consequence of the maximum principle.

Lemma 6.2. *Suppose that the initial metric g_0 satisfies $0 < \psi_s \leq 1$ and that $\lim_{s \rightarrow 0^+} \psi_s = 1$ and $\lim_{s \nearrow \infty} \psi_s = 0$ under the Ricci flow, then $0 < \psi_s \leq 1$ for as long as the solution to Ricci flow exists.*

Proof. Denoting $v = \psi_s$, then by [1, Equation (16)] the evolution of v is

$$v_t = v_{ss} + \frac{n-2}{\psi} v v_s + \frac{n-1}{\psi^2} (1-v^2)v.$$

Since $\lim_{s \nearrow \infty} \psi_s = 0$ under the flow, given any $\varepsilon > 0$, there exists $R = R(\varepsilon)$ such that $|v| < \varepsilon$ for $s \in [R, \infty)$. Define $w := e^{-\mu t} v + 2\varepsilon$, where $\mu > 0$ is to be specified, then $w > \varepsilon$ for all $s \geq 0$. We claim that $w > 0$ for as long as the solution exists. Suppose not, then there exists a first interior space-time point (t_0, s_0) where w achieves zero and such that at (t_0, s_0) ,

$$v = -2\varepsilon e^{\mu t}, \quad v_s = 0, \quad v_{ss} \geq 0, \quad w_t \leq 0.$$

and

$$\begin{aligned} 0 &\geq e^{\mu t} w_t|_{(t_0, s_0)} = v_t - \mu v \\ &\geq \frac{n-1}{\psi^2} (1-v^2)v - \mu v \\ &= -2\varepsilon e^{-\mu t} \left(\frac{n-1}{\psi^2} (1 - \varepsilon^2 e^{-2\mu t}) - \mu \right) \\ &> -2\varepsilon e^{-\mu t} \left(\frac{n-1}{\psi^2} - \mu \right) \\ &> 0 \end{aligned}$$

if we pick μ large enough. However, $0 \geq e^{\mu t} w_t|_{(t_0, s_0)} > 0$ is a contradiction. So the claim is proved. Since $\varepsilon > 0$ is arbitrary, we have $v > 0$. That $v \leq 1$ also follows from the maximum principle, so the lemma is proved. \square

Remark 6.3. The condition $\psi_s > 0$ can be interpreted as the absence of minimal sphere in the manifold, cf. [16, 25].

We now prove the main results in this paper.

Proof of Theorem 1.1. Let $n + 1 \geq 3$ and fix $\lambda > 0$. By patching formal solutions and smoothing it, there is a family of smooth profile functions at $\tau = \tau_0$ such that each such function z_0 satisfies $0 < z^-(u, \tau_0) < z_0 < z^+(u, \tau_0) < 1$ for $0 < u < \infty$. By Lemma 6.1, z_0 determines a complete rotationally symmetric metric g_0 on \mathbb{R}^{n+1} . It is straightforward to check that g_0 has bounded sectional curvatures everywhere, and K and L decay to zero at spatial infinity. In particular, we choose $\psi_s > 0$ initially, so (\mathbb{R}^{n+1}, g_0) does not contain any minimal sphere. Then exists unique solution $g(t)$ to Ricci flow starting from g_0 [8, 26]. The manifold $(\mathbb{R}^{n+1}, g(t))$ does not contain any minimal sphere by Lemma 6.2. Moreover, $g(t)$ is immortal [16].

The profile $z(u, \tau)$ of $g(t)$ is the unique solution of equation (3.1) for $0 < u < \infty$ and $\tau \geq \tau_0$, with boundary condition $z(0, \tau) = 1$, asymptotic condition $\lim_{u \nearrow \infty} z(u, \tau) = 0$, and initial data $z(u, \tau_0) = z_0$. By the comparison principle (Proposition 5.6), $0 < z^-(u, \tau) \leq z(u, \tau) \leq z^+(u, \tau)$. In particular, $z(u, \tau)$ defines a complete, rotationally symmetric, smooth metric $g(t)$ on \mathbb{R}^{n+1} by Lemma 6.1.

As $t \nearrow \infty$, the asymptotic behaviour of the solution agrees with that of the barriers, and hence with that of the formal solution. In particular, the sectional curvatures of $K(t)$ and $L(t)$ of $g(t)$ achieve the maximum at the origin \mathcal{O} and

$$K(t)|_{\mathcal{O}} = L(t)|_{\mathcal{O}} \sim t^{\lambda-1}.$$

So part (1) of Theorem 1.1 is proved. Moreover, $z^- \leq z(u, \tau) \leq z^+$ for any $\tau < \infty$, and the solution $z(u, \tau)$ exhibits the asymptotic behaviour of z^\pm . Near the origin, $z(u, \tau)$ converges uniformly to the Bryant soliton profile function for $0 < u < R_D e^{-\lambda\tau}$. Near spatial infinity, i.e., as $u \nearrow \infty$, $z(u, \tau) \searrow 0$ at a rate depending on λ as is given in (3.11), and so the sectional curvatures K and L are asymptotically flat according to (3.12) and (3.13), respectively. Thus, $g(t)$ has the asymptotic behaviour described in parts (2) and (3) of Theorem 1.1.

Therefore, Theorem 1.1 is proved. \square

REFERENCES

- [1] Sigurd Angenent and Dan Knopf, *An example of neckpinching for Ricci flow on S^{n+1}* , Math. Res. Lett. **11** (2004), no. 4, 493–518.
- [2] Sigurd B. Angenent, James Isenberg, and Dan Knopf, *Degenerate neckpinches in Ricci flow*, J. Reine Angew. Math. **709** (2015), 81–117.
- [3] Alexander Appleton, *Curvature blow-up in doubly-warped product metrics evolving by Ricci flow*, available at <https://arxiv.org/abs/1903.09936>.
- [4] Christoph Böhm and Burkhard Wilking, *Manifolds with positive curvature operators are space forms*, Ann. of Math. (2) **167** (2008), no. 3, 1079–1097.
- [5] Simon Brendle, *Rotational symmetry of self-similar solutions to the Ricci flow*, Invent. Math. **194** (2013), no. 3, 731–764.

- [6] ———, *Rotational symmetry of Ricci solitons in higher dimensions*, J. Differential Geom. **97** (2014), no. 2, 191–214.
- [7] Robert L. Bryant, *Ricci flow solitons in dimension three with $SO(3)$ -symmetries*, available at <http://www.math.duke.edu/~bryant/3DRotSymRicciSolitons.pdf>.
- [8] Bing-Long Chen and Xi-Ping Zhu, *Uniqueness of the Ricci flow on complete noncompact manifolds*, J. Differential Geom. **74** (2006), no. 1, 119–154.
- [9] Bennett Chow, *The Ricci flow on the 2-sphere*, J. Differential Geom. **33** (1991), no. 2, 325–334.
- [10] Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni, *The Ricci flow: techniques and applications. Part I*, Mathematical Surveys and Monographs, vol. 135, American Mathematical Society, Providence, RI, 2007.
- [11] Bennett Chow, Peng Lu, and Lei Ni, *Hamilton's Ricci flow*, Graduate Studies in Mathematics, vol. 77, American Mathematical Society, Providence, RI; Science Press Beijing, New York, 2006.
- [12] Xianzhe Dai and Li Ma, *Mass under the Ricci flow*, Comm. Math. Phys. **274** (2007), no. 1, 65–80. MR2318848
- [13] P. Daskalopoulos and Manuel del Pino, *Type II collapsing of maximal solutions to the Ricci flow in \mathbb{R}^2* , Ann. Inst. H. Poincaré Anal. Non Linéaire **24** (2007), no. 6, 851–874.
- [14] P. Daskalopoulos and R. Hamilton, *Geometric estimates for the logarithmic fast diffusion equation*, Comm. Anal. Geom. **12** (2004), no. 1-2, 143–164.
- [15] Panagiota Daskalopoulos and Natasa Sesum, *Type II extinction profile of maximal solutions to the Ricci flow in \mathbb{R}^2* , J. Geom. Anal. **20** (2010), no. 3, 565–591.
- [16] Francesco Di Giovanni, *Rotationally symmetric Ricci flow on \mathbb{R}^{n+1}* , available at <https://arxiv.org/abs/1904.09555>.
- [17] David Garfinkle and James Isenberg, *The modeling of degenerate neck pinch singularities in Ricci flow by Bryant solitons*, J. Math. Phys. **49** (2008), no. 7, 073505, 10.
- [18] Hui-Ling Gu and Xi-Ping Zhu, *The existence of type II singularities for the Ricci flow on S^{n+1}* , Comm. Anal. Geom. **16** (2008), no. 3, 467–494.
- [19] Richard S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom. **17** (1982), no. 2, 255–306.
- [20] ———, *Four-manifolds with positive curvature operator*, J. Differential Geom. **24** (1986), no. 2, 153–179.
- [21] James Isenberg and Haotian Wu, *Mean curvature flow of noncompact hypersurfaces with Type-II curvature blow-up*, J. Reine Angew. Math. **754** (2019), 225–251.
- [22] James Isenberg, Haotian Wu, and Zhou Zhang, *Mean curvature flow of noncompact hypersurfaces with Type-II curvature blow-up. II*, Preprint.
- [23] ———, *Precise asymptotics of type-iiib solutions to mean curvature flow*, Preprint.
- [24] Yan Li, Gang Tian, and Xiaohua Zhu, *Singular limits of Kähler-Ricci flow on Fano G -manifolds*, available at <https://arxiv.org/abs/1807.09167>.
- [25] Todd A. Oliynyk and Eric Woolgar, *Rotationally symmetric Ricci flow on asymptotically flat manifolds*, Comm. Anal. Geom. **15** (2007), no. 3, 535–568.
- [26] Wan-Xiong Shi, *Deforming the metric on complete Riemannian manifolds*, J. Differential Geom. **30** (1989), no. 1, 223–301.
- [27] Maxwell Stolarski, *Eguchi-Hanson singularities in $U(2)$ -invariant Ricci flow*, available at <https://arxiv.org/abs/1905.00087v1>.
- [28] Valentino Tosatti and Yuguang Zhang, *Infinite-time singularities of the Kähler-Ricci flow*, Geom. Topol. **19** (2015), no. 5, 2925–2948.
- [29] Haotian Wu, *On Type-II singularities in Ricci flow on \mathbb{R}^N* , Comm. Partial Differential Equations **39** (2014), no. 11, 2064–2090.

- [30] Shing-Tung Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I*, Comm. Pure Appl. Math. **31** (1978), no. 3, 339–411.

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, NSW
2006, AUSTRALIA

Email address: `haotian.wu@sydney.edu.au`