

Discrete Discreet Asymptotics

Nalini Joshi

@monsoon0



Supported by the London Mathematical Society and the Australian Research Council

Divergent Series



Divergent Series

- From Littlewood's preface to GH Hardy's *Divergent Series*



Divergent Series

- From Littlewood's preface to GH Hardy's *Divergent Series*

“Abel wrote in 1828: ‘Divergent series are the invention of the devil, and it is shameful to base on them any demonstration whatsoever.’”



Divergent Series

- From Littlewood's preface to GH Hardy's *Divergent Series*

“Abel wrote in 1828: ‘Divergent series are the invention of the devil, and it is shameful to base on them any demonstration whatsoever.’”

“Then came a time when it was found that something after all could be done about them. This is now a matter of course, but in the early years of the century the subject, while in no way mystical or unrigorous, was regarded as sensational, and about the present title, now colourless, there hung an aroma of paradox and audacity.”



Series

$$1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots$$


$$1 - 1 + 1 - 1 + 1 - 1 + \dots$$

$$1 - \epsilon + \epsilon^2 - \epsilon^3 + \epsilon^4 + \dots$$

$$1 + \epsilon + 2!\epsilon^2 + 3!\epsilon^3 + \dots$$

Series

$$1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots$$


$$1 - 1 + 1 - 1 + 1 - 1 + \dots$$

$$1 - \epsilon + \epsilon^2 - \epsilon^3 + \epsilon^4 + \dots$$

$$1 + \epsilon + 2!\epsilon^2 + 3!\epsilon^3 + \dots$$

Series

$$1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots$$

$$1 - 1 + 1 - 1 + 1 - 1 + \dots$$

$$1 - \epsilon + \epsilon^2 - \epsilon^3 + \epsilon^4 + \dots$$

$$1 + \epsilon + 2!\epsilon^2 + 3!\epsilon^3 + \dots$$

Interpretation?

$$\frac{1}{1 - 2x} = 1 + 2x + 4x^2 + 8x^3 + \dots$$

$$x = 1$$

\Downarrow

$$-1 = 1 + 2 + 4 + 8 + \dots$$

GH Hardy, *Divergent Series*, 1949

Asymptotics

Asymptotics is the approximation of functions in limits.

$$\text{As } x \underset{\gamma}{\rightarrow} x_0$$

Asymptotics

Asymptotics is the approximation of functions in limits.

$$\text{As } x \underset{\gamma}{\rightarrow} x_0$$

$$f(x) \ll g(x) \iff \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0$$

Asymptotics

Asymptotics is the approximation of functions in limits.

$$\text{As } x \xrightarrow[\gamma]{} x_0$$

$$f(x) \ll g(x) \iff \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0$$

$$f(x) \sim g(x) \iff \lim_{x \rightarrow x_0} \frac{f(x) - g(x)}{g(x)} = 0$$

Asymptotics

Asymptotics is the approximation of functions in limits.

$$\text{As } x \xrightarrow[\gamma]{} x_0$$

$$f(x) \ll g(x) \iff \lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0$$

$$f(x) \sim g(x) \iff \lim_{x \rightarrow x_0} \frac{f(x) - g(x)}{g(x)} = 0$$

$$f(x) = \mathcal{O}(g(x)) \iff \exists M \text{ s.t. } \left| \frac{f(x)}{g(x)} \right| < M$$

$$\forall x \in \gamma$$

Asymptotic Series

- f is said to be asymptotic to a series $\sum_{n=0}^{\infty} a_n (x - x_0)^n$
iff $\forall N \in \mathbb{N}$

$$\lim_{x \rightarrow x_0} \frac{f(x) - \sum_{n=0}^N a_n (x - x_0)^n}{(x - x_0)^N} = 0$$

and we write

$$f(x) \sim \sum_{n=0}^{\infty} a_n (x - x_0)^n, \text{ as } x \rightarrow x_0$$

Asymptotic Series

- f is said to be asymptotic to a series $\sum_{n=0}^{\infty} a_n (x - x_0)^n$
iff $\forall N \in \mathbb{N}$

$$\lim_{x \rightarrow x_0} \frac{f(x) - \sum_{n=0}^N a_n (x - x_0)^n}{(x - x_0)^N} = 0$$

and we write

$$f(x) \sim \sum_{n=0}^{\infty} a_n (x - x_0)^n, \text{ as } x \rightarrow x_0$$

- The series is not required to converge.

An example

$$u'' - u' \left(1 - \frac{1}{x}\right) = 0$$

- One solution is

$$Ei(x) = \int_{-\infty}^x \frac{e^t}{t} dt$$
$$\sim \frac{e^x}{x} \sum_{n=0}^{\infty} \frac{n!}{x^n}, \quad x \rightarrow +\infty$$

- Other solutions differ from this one by a constant.

$$Ei(x) + C$$

An example

$$u'' - u' \left(1 - \frac{1}{x} \right) = 0$$

- One solution is

$$Ei(x) = \int_{-\infty}^x \frac{e^t}{t} dt$$

$$\sim \frac{e^x}{x} \sum_{n=0}^{\infty} \frac{n!}{x^n}, \quad x \rightarrow +\infty$$

• *Divergent*

- Other solutions differ from this one by a constant.

$$Ei(x) + C$$

Approximation

- Consider $F(x) = x e^{-x} Ei(x)$

$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} \text{ as } x \rightarrow \infty \quad \Leftrightarrow \quad \forall N \quad \left| \frac{F(x) - \sum_{k=0}^N k! / x^k}{N! / x^N} \right| \rightarrow 0$$

Approximation

- Consider $F(x) = x e^{-x} Ei(x)$

$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} \text{ as } x \rightarrow \infty \quad \Leftrightarrow \quad \forall N \left| \frac{F(x) - \sum_{k=0}^N k! / x^k}{N! / x^N} \right| \rightarrow 0$$

$$\sum_{n=0}^{10} \frac{n!}{100^n} = 1.0102062527748311680 \dots$$

Approximation

- Consider $F(x) = x e^{-x} Ei(x)$

$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} \text{ as } x \rightarrow \infty \quad \Leftrightarrow \quad \forall N \left| \frac{F(x) - \sum_{k=0}^N k! / x^k}{N! / x^N} \right| \rightarrow 0$$

$$\sum_{n=0}^{10} \frac{n!}{100^n} = 1.0102062527748311680 \dots$$

$$\sum_{n=0}^{100} \frac{n!}{100^n} = 1.0102062527748357112 \dots$$

Approximation

- Consider $F(x) = x e^{-x} Ei(x)$


$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} \text{ as } x \rightarrow \infty \quad \Leftrightarrow \quad \forall N \left| \frac{F(x) - \sum_{k=0}^N k!/x^k}{N!/x^N} \right| \rightarrow 0$$

$$\sum_{n=0}^{10} \frac{n!}{100^n} = 1.0102062527748311680 \dots$$

$$\sum_{n=0}^{100} \frac{n!}{100^n} = 1.0102062527748357112 \dots$$

$$\sum_{n=0}^{281} \frac{n!}{100^n} = 732496.06921461904157 \dots$$


Truncation at least term



- Truncate at the least term

$$F(x) = \sum_{n=0}^{N-1} \frac{n!}{x^n} + R_N$$

Truncation at least term



- Truncate at the least term

$$F(x) = \sum_{n=0}^{N-1} \frac{n!}{x^n} + R_N$$

- *Exponentially small*
- *N depends on x*

The identification problem

- How do we distinguish

$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n}$$

$$\tilde{F}(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} + \frac{e^{-x}C}{x}$$

in the limit $x \rightarrow +\infty$?

- E.g., if $C=1$


$$\frac{e^{-100}}{100} = 3.7200759760208359630 \times 10^{-46}$$

The identification problem

- How do we distinguish

- *Discreetly* hidden beyond all orders

$$F(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n}$$

$$\tilde{F}(x) \sim \sum_{n=0}^{\infty} \frac{n!}{x^n} + \frac{e^{-x}C}{x}$$


in the limit $x \rightarrow +\infty$?


- E.g., if $C=1$

$$\frac{e^{-100}}{100} = 3.7200759760208359630 \times 10^{-46}$$

Why should we care?

Chaos

- “Does the flap of a butterfly’s wings in Brazil set off a tornado in Texas?”
- To forecast future behaviour, we need to know initial states with infinite precision.
- This has become synonymous with *chaos*, but is also present in ordered, non-chaotic systems.



TC Gavin 1997 SW Pacific

Order

- The Painlevé equations, which arise as reductions of soliton equations

$$w_{\tau} + 6 w w_{\xi} + w_{\xi\xi\xi} = 0$$

$$\begin{cases} w = -2 y(x) - 2 \tau \\ x = \xi + 6 \tau^2 \end{cases}$$

$$\Rightarrow \begin{cases} w_{\tau} &= -24 \tau y_x - 2 \\ w_{\xi} &= -2 y_x \\ w_{\xi\xi\xi} &= -2 y_{xxx} \end{cases}$$


Order

- The Painlevé equations, which arise as reductions of soliton equations

$$w_{\tau} + 6 w w_{\xi} + w_{\xi\xi\xi} = 0$$

$$\begin{cases} w = -2 y(x) - 2 \tau \\ x = \xi + 6 \tau^2 \end{cases}$$

$$\Rightarrow \begin{cases} w_{\tau} &= -24 \tau y_x - 2 \\ w_{\xi} &= -2 y_x \\ w_{\xi\xi\xi} &= -2 y_{xxx} \end{cases}$$


$$y'' = 6 y^2 - x$$

Applications

- Electrical structures of interfaces in steady electrolysis *L. Bass, Trans Faraday Soc 60 (1964) 1656–1663*
- Spin-spin correlation functions for the 2D Ising model *TT Wu, BM McCoy, CA Tracy, E Barouch Phys Rev B13 (1976) 316–374*
- Spherical electric probe in a continuum gas *PCT de Boer, GSS Ludford, Plasma Phys 17 (1975) 29–41*
- Cylindrical Waves in General Relativity *S Chandrasekar, Proc. R. Soc. Lond. A 408 (1986) 209–232*
- Non-perturbative 2D quantum gravity *Gross & Migdal PRL 64(1990) 127-130*
- Orthogonal polynomials with non-classical weight function *AP Magnus J. Comput Appl. Anal. 57 (1995) 215–237*
- Level spacing distributions and the Airy kernel *CA Tracy, H Widom CMP 159 (1994) 151–174*
- Spatially dependent ecological models: *Joshi & Morrison Anal Appl 6 (2008) 371-381*
- Gradient catastrophe in fluids: *Dubrovin, Grava & Klein J. Nonlin. Sci 19 (2009) 57-94*


What do we know about the solutions of these equations?



$$u(0) = 0, \quad u'(0) = 0$$

General Solutions

- Movable poles
- Transcendentality of general solutions
- Special solutions
- Asymptotic behaviours



Fornberg & Weideman 2009

P_I

Tronquée Solutions





FIG. 3.1. Magnitude of the solution $u(z)$ to the P_I equation in case of ICs $u(0) = -0.1875$, $u'(0) = 0.3049$, displayed over the domain $z = x + iy$, $-10 \leq x \leq 10$, $-10 \leq y \leq 10$.

Real Solutions

Consider $P_1 \quad w_{tt} = 6w^2 - t$ for $w(t), t \in \mathbb{R}$



The Real Tritronquée


- Theorem: \exists **unique** solution $Y(x)$ of PI which has asymptotic expansion

$$y_f = -\sqrt{\frac{x}{6}} \sum_{k=0}^{\infty} \frac{a_k}{(x^{1/2})^{5k}}, \text{ in } |\arg(x)| \leq 4\pi/5$$

and

- $Y(x)$ is real for real x
- Its interval of existence I contains \mathbb{R}
- $Y(x)$ lies below Π .
- It is monotonically decaying in I .

Exponential Terms



What about global dynamics?

Perturbed Form

- In Boutroux's coordinates:

$$w_1 = t^{1/2} u_1(z), \quad w_2 = t^{3/4} u_2(z), \quad z = \frac{4}{5} t^{5/4}$$

$$\begin{pmatrix} \dot{u}_1 \\ \dot{u}_2 \end{pmatrix} = \begin{pmatrix} u_2 \\ 6u_1^2 - 1 \end{pmatrix} - \frac{1}{5z} \begin{pmatrix} 2u_1 \\ 3u_2 \end{pmatrix}$$

- a perturbation of an elliptic curve as $|z| \rightarrow \infty$

$$E = \frac{u_2^2}{2} - 2u_1^3 + u_1 \quad \Rightarrow \quad \frac{dE}{dz} = \frac{1}{5z} (6E + 4u_1)$$

Perturbed Form

- In Boutroux's coordinates:

$$w_1 = t^{1/2} u_1(z), \quad w_2 = t^{3/4} u_2(z), \quad z = \frac{4}{5} t^{5/4}$$

$$\begin{pmatrix} \dot{u}_1 \\ \dot{u}_2 \end{pmatrix} = \begin{pmatrix} u_2 \\ 6u_1^2 - 1 \end{pmatrix} - \frac{1}{5z} \begin{pmatrix} 2u_1 \\ 3u_2 \end{pmatrix}$$

- a perturbation of an elliptic curve as $|z| \rightarrow \infty$

$$E = \frac{u_2^2}{2} - 2u_1^3 + u_1 \quad \Rightarrow \quad \frac{dE}{dz} = \frac{1}{5z} (6E + 4u_1)$$

Projective Space

- What if x, y become unbounded?
- Use projective geometry: $x = \frac{u}{w}, y = \frac{v}{w}$
 $[x, y, 1] = [u, v, w] \in \mathbb{CP}^2$


- The level curves of P_I are now

$$F_I = wv^2 - 4u^3 + g_2uw^2 + g_3w^3$$

all intersecting at the **base point** $[0, 1, 0]$.

\Rightarrow To describe solutions, **resolve** the flow through this point

Resolving a base pt



Resolution

- “Blow up” the singularity or base point:

$$f(x, y) = y^2 - x^3$$

$$(x, y) = (x_1, x_1 y_1)$$


$$\Rightarrow x_1^2 y_1^2 - x_1^3 = 0$$

$$\Leftrightarrow x_1^2 (y_1^2 - x_1) = 0$$

- Note that

$$x_1 = x, y_1 = y/x$$


$$y^2 = x^3$$



Method

$$f(x, y) = y^2 - x^3$$


$$(x, y) = (x_1, x_1 y_1)$$



$$f(x_1, x_1 y_1) = x_1^2 (y_1^2 - x_1)$$

$$L_1 : \textcircled{-1}$$


$$y_1^2 = x_1$$




$$f_1(x_1, y_1) = y_1^2 - x_1$$

$$f_1(x_2, y_2) = y_2(y_2 - x_2)$$

$$(x_1, y_1) = (x_2, y_2)$$




$$(x_2, y_2) = (x_3, x_3 y_3)$$



$$f_2(x_2, y_2) = y_2 - x_2$$

$$f_2(x_3, x_3 y_3) = x_3(y_3 - 1)$$

Initial-Value Space




$$L_1 : x_1 = 0$$

$$L_2 : y_2 = 0$$

$$L_3 : x_3 = 0$$

Now the space is compactified and regularised.


Initial-Value Space



Now the space is compactified and regularised.

Unifying Property


The space of initial values of
a Painlevé system is
resolved by “blowing up” 9
points in \mathbb{CP}^2
(or 8 points in $\mathbb{P}^1 \times \mathbb{P}^1$)



Okamoto 1979

Sakai 2001


Sakai's Description I



Initial-value spaces of all continuous and discrete Painlevé equations


P_I

L_9



P_1


L_9



P_1


L_9

$E_8^{(1)}$




P_I



L_9 autonomous eqn





dP_I



dP_I





dP_1



degenerate autonomous limit

dP_1



qP_1
 $h_v - e_6$

$\mathbb{P}^1 \times \mathbb{P}^1$

$h_u - e_3$

$h_u - e_2 - e_6$

$h_v - e_1 - e_3$

v

u

e_8

e_6

e_7


e_2

e_3

e_4

e_5

e_1



qP_1
 $h_v - e_6$


$\mathbb{P}^1 \times \mathbb{P}^1$

$h_u - e_3$

$h_u - e_2 - e_6$

$h_v - e_1 - e_3$


$A_7^{(I)}$



qP_1
 $h_v - e_6$

$\mathbb{P}^1 \times \mathbb{P}^1$

$h_u - e_3$



Symmetric dP1


$$w_{n+1} + w_n + w_{n-1} = \frac{\alpha n + \beta}{w_n} + \gamma$$

- Consider $n \rightarrow \infty$
- General behaviours are close to elliptic functions
- Special solutions are given by power series

Joshi 1997

Vereschagin 1995

Solutions



Solution orbits of scalar dP1 on the Riemann sphere (where the north pole is infinity).

Scaling

$$\begin{cases} w_{2k} &= \frac{u(s)}{\epsilon^{1/2}} \\ w_{2k-1} &= \frac{v(s)}{\epsilon^{1/2}} \end{cases} \quad s = \epsilon n$$

- dPI becomes

$$(v(s + \epsilon) + u(s) + v(s - \epsilon))u(s) = \alpha s + \epsilon \beta + \epsilon^{1/2} \gamma u(s)$$

$$(u(s + \epsilon) + v(s) + u(s - \epsilon))v(s) = \alpha s + \epsilon \beta + \epsilon^{1/2} \gamma u(s)$$

- Series expansions as $\epsilon \rightarrow 0$

$$u(s) \sim \sum_{m=0}^{\infty} \epsilon^{m/2} u_m(s)$$

$$v(s) \sim \sum_{m=0}^{\infty} \epsilon^{m/2} v_m(s)$$

Types of solutions

- Type A

$$u \sim \pm \sqrt{-\alpha s} + \frac{\gamma \epsilon^{1/2}}{2} \mp \frac{(4\beta - \gamma^2)\epsilon}{8\sqrt{-\alpha s}} + \dots$$

$$v \sim \mp \sqrt{-\alpha s} + \frac{\gamma \epsilon^{1/2}}{2} \pm \frac{(4\beta - \gamma^2)\epsilon}{8\sqrt{-\alpha s}} + \dots$$

- Type B

$$u = v \sim \pm \sqrt{\frac{\alpha s}{3}} + \frac{\gamma \epsilon^{1/2}}{6} \mp \pm \frac{\sqrt{3}(12\beta + \gamma^2)\epsilon}{72\sqrt{\alpha s}} + \dots$$


Late-order terms: Type A

$$u_m \sim \frac{\Lambda_1 \Gamma\left(\frac{m-1}{2}\right)}{(i\pi s/2)^{\frac{m-1}{2}}} + \frac{\Lambda_2 \Gamma\left(\frac{m-1}{2}\right)}{(-i\pi s/2)^{\frac{m-1}{2}}}$$
$$v_m \sim \frac{\Lambda_3 \Gamma\left(\frac{m-1}{2}\right)}{(i\pi s/2)^{\frac{m-1}{2}}} + \frac{\Lambda_4 \Gamma\left(\frac{m-1}{2}\right)}{(-i\pi s/2)^{\frac{m-1}{2}}}$$

- Optimal truncation

$$u(s) \sim \sum_{m=0}^{N_o} \epsilon^{m/2} u_m(s) + S_1 \Lambda_1 (-i)^{s/\epsilon} + S_2 \Lambda_2 i^{s/\epsilon}$$

Stokes Sectors: Type A



Joshi & Lustri 2015

What about q -discrete Painlevé equations?

qP1

$$\Rightarrow \quad \overline{w} \, \underline{w} = \frac{1}{w} - \frac{1}{\xi w^2} \quad (\text{qP}_I)$$

$$\overline{w} = w(q \xi), \quad w = w(\xi), \quad \underline{w} = w(\xi/q)$$

- A limiting form of qP3, rescaled

$$\overline{g} \, \underline{g} = \frac{\alpha x}{g} + \frac{\beta}{g^2} \quad \text{Ramani \& Grammaticos (1996)}$$

$$\overline{g} = g(\tilde{q} x), \quad \underline{g} = g(x/\tilde{q})$$

$$\mapsto \text{PI:} \quad y'' = 6 y^2 - t \quad \text{in continuum limit.}$$

Behaviours near fixed points

$$\overline{w} \sim w, \quad \underline{w} \sim w, \quad |\xi| \rightarrow \infty$$

$$\Rightarrow w^4 = w + \mathcal{O}(1/\xi)$$

$$\Rightarrow w = \begin{cases} \omega + \mathcal{O}(1/\xi) \\ \mathcal{O}(1/\xi) \end{cases} \quad \omega^3 = 1$$

- qP_I is invariant under rotation by argument $2\pi/3$, so ω can be replaced by unity.
- The second case lies in neighbourhood of a merger of two base points: $(1/\xi, 0)$, $(q/\xi, 0)$.

Near unity

- Near $w = 1$, $\underline{w} = 1$, \exists a formal series solution

$$w = \sum_{n=0}^{\infty} a_n t^n, \quad a_0 = 1$$

where

$$\begin{aligned} & a_n (q^n + 1 + q^{-n}) \\ &= - \sum_{l=1}^{n-1} a_l a_{n-l} (q^{(2l-n)} + 1) \\ &\quad - \sum_{m=1}^{n-1} \sum_{j=0}^{n-m} \sum_{l=0}^m a_j a_{n-m-j} a_l a_{m-l} q^{(n-m-2j)} \end{aligned}$$

Near zero

- Near $w = 1/\xi$, $\underline{w} = q/\xi$, \exists a formal series solution

where

$$w(\xi) = \sum_{n=1}^{\infty} \frac{b_n}{\xi^n}$$

$$b_1 = 1, \quad b_2 = 0, \quad b_3 = 0$$

$$b_n = \sum_{r=2}^{n-2} \sum_{k=1}^{r-1} \sum_{m=1}^{n-r-1} b_k b_{r-k} b_m b_{n-r-m} q^{(r-2k)}, \quad n \geq 4$$

Step 1: divergence

The coefficients of the asymptotic series grow very fast:

$$b_{3p+1} \underset{p \rightarrow \infty}{=} \mathcal{O} \left(|q|^{3p(p-1)/2} \prod_{k=0}^{p-1} (1 + q^{-3k})^2 \right), |q| > 1$$

$$b_{3p+2} = 0, \quad b_{3p+3} = 0, \quad \forall p \geq 0$$

Step 2: Analytic Sum

- Use of the Borel-Ritt theorem provides an analytic function W s.t.

$$W(\xi) \sim \sum_{n=1}^{\infty} \frac{b_n}{\xi^n}$$

Step 3: Linearisation

- The linearisation around W satisfies

$$\overline{P} + \left(2 \frac{\overline{W}}{W} - \frac{1}{W^2 \underline{W}} \right) P + \frac{\overline{W}}{\underline{W}} \underline{P} = 0$$

which has solutions with behaviours

$$P^{\pm}(\xi) \sim q^{\pm 3n(n \mp 5/3)/2}$$

$$\xi = \xi_0 q^n$$

Step 4: True Solutions

- The perturbed q-difference equation gives

$$v_n = \beta_0 P_n - P_n \sum_{j=n}^{n_0-1} \frac{W_j W_{j-1}}{P_j P_{j-1}} \sum_{k=k_0}^{j-1} \frac{P_k \mathcal{R}_2(v_k, v_{k-1}, t_k)}{W_{k+1} W_k}$$

where

$$\|\mathcal{R}_2(v, \underline{v}, t)\| \leq C_1 \|\mathbf{v}\|^2 + C_2 |t|$$

$$\|\nabla \mathcal{R}_2\| \leq C_3 \|\mathbf{v}\| + C_4 |t|$$

- The contraction mapping theorem provides a true solution.

Quicksilver solution

- The vanishing solution approaches two base points.
- Its series expansion is divergent.
- We prove a true solution exists with this behaviour; it does not lie on a singularity of the underlying elliptic curve. So it is different to the *tritronquée* solutions of the Painlevé equations.

⇒ new name: *quicksilver* solution

- It is unstable in initial-value space.

Joshi, Stud Appl Math (2014)

Comparison

PI


- No rational or classical solutions
- Leading-order behaviour is elliptic
- Two types of solutions described by asymptotic behaviours
- Tronquée solutions are asymptotic to a power series in a large sector

qPI

- No algebraic or solutions in terms of linear eqns *Nishioka (2010)*
- Leading-order behaviour is elliptic
- Four types of solutions described by asymptotic behaviours *J (2014)*
- Quasi-stationary solutions are asymptotic to a power series in a large region *J (2014)*

Summary

- New mathematical models of physics pose new questions for applied mathematics
- **Global** dynamics of solutions of non-linear equations, whether they are differential or discrete, can be found through geometry.
- Geometry provides the only **analytic approach** available in \mathbb{C} for discrete equations.
- Tantalising questions about **finite properties** of solutions remain open.



The mathematician's pattern's, like those of the painter's or the poet's, must be beautiful, the ideas, like the colours or the words, must fit together in a harmonious way. *GH Hardy, A Mathematician's Apology, 1940*