

ON FARBER'S INVARIANTS FOR SIMPLE $2q$ -KNOTS

JONATHAN A. HILLMAN

ABSTRACT. Let K be a simple $2q$ -knot with exterior X . We show directly how the Farber quintuple $(A, \Pi, \alpha, \ell, \psi)$ determines the homotopy type of X if the torsion subgroup of $A = \pi_q(X)$ has odd order. We comment briefly on the possible role of the EHP sequence in recovering the boundary inclusion from the duality pairings ℓ and ψ . Finally we reformulate the Farber quintuple as an hermitean self-duality of an object in an additive category with involution.

In a series of papers in the early 1980s Farber showed that stable high dimensional knots could be classified in terms of stable homotopy theory, and in particular that if $q \geq 4$ simple $2q$ -knots may be classified up to isotopy by quintuples $(A, \Pi, \alpha, \ell, \psi)$, where A, Π are $\mathbb{Z}[Z]$ -modules, α is a monomorphism and ℓ and ψ are sesquilinear pairings [6]. Certain significant special cases were dealt with by Kearton [14] and Kojima [15].

Farber's argument for showing that his invariant determined the knot involves Spanier-Whitehead duality for highly connected Seifert hypersurfaces and Wall's embedding theorem (to interpolate between different choices of such hypersurfaces). An alternative approach (avoiding Seifert arguments) is suggested by the work of Lashof and Shaneson, who used surgery over $\pi_1 = Z$ to show that the exterior X of a high dimensional knot with group Z is determined up to homeomorphism by the homotopy type of the pair $(X, \partial X)$ [16]. There remains a possible ambiguity of order 2, and a further argument is needed to show that the exterior determines the knot if $(X, \partial X)$ is highly connected [20].

In this note we shall provide some evidence for the feasibility of such a homotopy-theoretic approach. We show first that the homotopy type of X is determined by the modules $A = \pi_q(X)$ and $\pi_{q+1}(X)$ and a k -invariant $\kappa(X)$. We then show that these invariants may be derived algebraically from the Farber quintuple. (In the case of $\kappa(X)$ we need to assume that the \mathbb{Z} -torsion submodule $T(A)$ has odd order.) The work of Farber implies that the duality pairings ℓ and ψ determine the inclusion of the boundary. We have not yet been able to verify this independently. However the longitude (i.e., the lift of the boundary inclusion to infinite cyclic covers) has trivial suspension, and the EHP sequence of Chapter XII of [25] suggests a close connection between the longitude and these duality pairings. Further progress appears to depend on having a more direct construction of the pairing ψ (or at least of the version Ψ used below for the case with $T(A)$ odd).

Finally, Quebbemann, Scharlau and Schulte have codified the notion of hermitean pairing in an additive category \mathcal{C} with an involution $*$ [19], and we shall show that Farber's invariant may be described in such terms. This approach has

1991 *Mathematics Subject Classification.* 57Q45.

Key words and phrases. EHP sequence, Farber quintuple, F -form, hermitean duality, simple knot.

proven useful in studying the factorization of some nontrivial classes of higher-dimensional knots into sums of indecomposable knots [8, 10]. It is also of interest in connection with criteria for a knot to be doubly null concordant [7].

1. KNOTS

An n -knot is a locally flat embedding $K : S^n \rightarrow S^{n+2}$. We shall assume that S^n has the standard orientation as the boundary of the unit ball in R^{n+1} , for all $n > 0$. The exterior of K is $X = X(K) = S^{n+2} - N(K)$, where $N(K) \cong S^n \times R^2$ is an open regular neighbourhood of the image of K . The orientations of the spheres determine an orientation for X and an isotopy class of homeomorphisms $S^n \times S^1 \cong \partial X$. The inclusion of S^1 into X determined by the *meridian* $* \times S^1 \subset \partial X$ induces an isomorphism on integral homology. In particular, there is a canonical generator t for $H = H_1(X(K); \mathbb{Z}) \cong \mathbb{Z}$, and so we may identify $\mathbb{Z}[H]$ with $\mathbb{Z}[Z] = \Lambda = \mathbb{Z}[t, t^{-1}]$.

Reattaching $S^n \times D^2$ to $X = X(K)$ along $\partial X \cong S^n \times S^1$ using the nontrivial twist map of $S^n \times S^1$ gives a homotopy $(n+2)$ -sphere. The embedding of $S^n \times \{0\}$ in this sphere gives an n -knot K^* , called the *Gluck reconstruction* of K , and any knot with exterior homeomorphic to X is isotopic to K or to K^* (up to change of orientations).

Let $p : X' \rightarrow X$ be the infinite cyclic cover space corresponding to the commutator subgroup π' . The modules $H_*(X; \Lambda) = H_*(X'; \mathbb{Z})$ are finitely generated, since X is compact and Λ is noetherian. Moreover $t - 1$ acts invertibly on $H_*(X; \Lambda)$, since X is an homology circle, and so these are torsion modules. The *longitude* $\lambda_K \in \pi_n(X)$ is the class of the inclusion $S^n \times \{1\} \rightarrow S^n \times S^1 \cong \partial X \subset X$. If V is a Seifert hypersurface for K then λ_K factors through the inclusion of $\partial V \cong S^n$ into V . Let $k : V/\partial V \rightarrow S^{n+1}$ be the cofibre of the natural map from V to $V/\partial V$. Then the Pontrjagin-Thom collapse of S^{n+2} onto $S(V/\partial V)$ provides a section to Sk . It follows easily that the suspension of λ_K is trivial [22].

The knot K is r -simple if $\pi_1(X) \cong \mathbb{Z}$ and $\pi_j(X) = 0$ for $1 < j \leq r$. The sum $K \# \tilde{K}$ of r -simple n -knots K and \tilde{K} is r -simple. If K is 1-simple then X is determined up to homeomorphism by the homotopy type of the pair $(X, \partial X)$ [16]. The knot K is *simple* if it is $[(n-1)/2]$ -simple and $n \geq 3$.

It is a consequence of Farber's work on stable knots that an r -simple n -knot K is determined by its exterior (i.e., $K \cong K^*$) if $3r > n$. In [20] Richter shows that if $3r \geq n$ and K is an r -simple n -knot the twist map of ∂X extends to a self homotopy equivalence of X and hence that K is determined by its complement, as a straightforward consequence of Williams' Poincaré Embedding Theorem. Richter's approach is close to the one taken here, in that he strives to work as far as possible in the homotopy category, before appealing to geometric topology through surgery. In particular, he subsequently gave a purely homotopy theoretic proof of William's theorem [21]. (The application to knot theory does however use the fact that K has a highly connected Seifert hypersurface, in an essential way.) Is there an ad hoc simplification for the case of simple knots?

2. MODULES AND EXTENSIONS

If M is a finitely generated left Λ -module let $T(M)$ be the maximal finite submodule of M . In the cases of interest here $T(M)$ is the \mathbb{Z} -torsion submodule of M . The quotient $M/T(M)$ is \mathbb{Z} -torsion free, and has projective dimension at most 1

($p.d._\Lambda M/T(M) \leq 1$). If R is a *right* Λ -module let \overline{R} denote the left module with the conjugate structure determined by $t.r = rt^{-1}$ for all $r \in R$.

The extension modules $Ext_\Lambda^i(M, \Lambda)$ are naturally right modules. Let $e^i M = \overline{Ext_\Lambda^i(M, \Lambda)}$. If M is a finite Λ -module then $e^0 M = e^1 M = 0$, and so the inclusion $T(M) \leq M$ induces isomorphisms $e^i M \cong e^i(M/T(M))$ for $i = 0, 1$ and $e^2 M \cong e^2 T(M)$. More generally, $Ext_\Lambda^2(M, N) \cong Ext_\Lambda^2(T(M), N)$, for any Λ -module N . If $p.d._\Lambda M = d$ then $Ext_\Lambda^d(M, N)$ and $Ext_\Lambda^d(M, \Lambda) \otimes_\Lambda N$ are naturally isomorphic, for if P_* is a finitely generated free resolution of M then $Hom_\Lambda(P_*, N) \cong Hom_\Lambda(P_*, \Lambda) \otimes_\Lambda N$, and $- \otimes_\Lambda N$ is right exact.

If P_* is a chain complex of finitely generated free left Λ -modules let $D(P)_j = \overline{Hom_\Lambda(P_{-j}, \Lambda)}$ be the dual complex, and let $P[n]_*$ be the n -fold suspension, with $P[n]_j = P_{j-n}$ for all j . Note that $D(D(P))_*$ is naturally isomorphic to P_* . If $H_*(P) = 0$ for $j \neq q$ or $q+1$ then P is determined up to chain homotopy equivalence by the modules $A = H_q(P_*)$ and $B = H_{q+1}(P_*)$ and a k -invariant $\kappa(P_*)$ in $Ext_\Lambda^2(A, B)$ [4]. This may be identified with the class of the sequence

$$0 \rightarrow B \rightarrow P_{q+1}/\partial P_{q+2} \rightarrow \text{Ker}(\partial_q) \rightarrow A \rightarrow 0$$

in $Ext_\Lambda^2(A, B) \cong Ext_\Lambda^2(T(A), B)$. Every such class is realizable by a free complex concentrated in degrees $[q, q+2]$.

A *simple Λ -complex concentrated in degrees $[q, q+2]$* is a finitely generated free Λ -complex C_* whose homology modules are Λ -torsion modules and are 0 except in degrees q and $q+1$. Let $A = H_q(C_*)$, $T = T(A)$ and $B = H_{q+1}(C_*)$ and let $k(A) \in Ext_\Lambda^1(A/T, T) \cong \overline{e^1 A} \otimes T$ and $\kappa(C_*) \in Ext_\Lambda^2(A, B) \cong \overline{e^2 T} \otimes B$ be the associated extension classes.

Theorem 1. *Let C_* be a simple Λ -chain complex concentrated in degrees $[q, q+2]$.*

- (1) *C is determined up to chain homotopy equivalence by the quintuple $(A/T(A), T(A), B, k(A), \kappa(C_*))$;*
- (2) *any such system of invariants with A and B finitely generated Λ -torsion modules and $p.d._\Lambda B \leq 1$ is realized by a simple Λ -complex;*
- (3) *$D_* = D(C)_*$ is a simple complex concentrated in degrees $[-q-2, -q]$, with invariants $(e^1 B, e^2 A, e^1 A, k_D, \kappa(D_*))$ where k_D and $\kappa(D_*)$ are determined by $\kappa(C_*)$ and $k(A)$, respectively.*

Proof. The module A is determined by $T = T(A)$, A/T and $k(A)$, and so the first assertion follows from the result of [4] cited above. Since A is a torsion module D_* is easily seen to be a simple Λ -complex. If $A_D = H_{-q-2}(D_*)$ and $B_D = H_{-q-1}(D_*)$ then $B_D \cong e^1 A$, $T_D = T(A_D) \cong e^2 A = e^2 T$ and $A_D/T_D \cong e^1 B$.

There are natural isomorphisms $T \cong e^2 e^2 A = e^2 T_D$ and $B \cong e^1 A_D$. Hence the transpositions $\tau_1 : \overline{e^1 A} \otimes T \cong \overline{T} \otimes e^1 A$ and $\tau_2 : \overline{e^2 T} \otimes B \cong \overline{B} \otimes e^2 T$ given by interchange of factors induce natural isomorphisms $e^1 A \otimes T \cong e^2 T_D \otimes B_D$ and $e^2 T \otimes B \cong e^1 A_D \otimes T_D$ between the abelian groups in which the extension classes for A and D_* and for C_* and A_D lie.

To verify (2) and see that the extension classes correspond as asserted we shall construct an explicit representative for the chain homotopy type determined by the invariants $(A/T, T, B, k(A), \kappa(C_*))$. We shall assume for simplicity of notation that $q = 0$. Let $P_1 \rightarrow P_0$, $Q_1 \rightarrow Q_0$ and $E_2 \rightarrow E_1 \rightarrow E_0$ be finite free resolutions of A/T , B and T , respectively, and let $k : P_1 \rightarrow E_0$ and $\kappa : E_2 \rightarrow Q_0$ be homomorphisms representing $k(A)$ and $\kappa(C_*)$, respectively. Then we may use a mapping cone

construction (twice) as in Satz 7.6 of [4] to construct a complex

$$(1) \quad E_2 \oplus Q_1 \xrightarrow{\Delta_2(\kappa)} P_1 \oplus E_1 \oplus Q_0 \xrightarrow{\Delta_1(k)} P_0 \oplus E_0$$

with these invariants. Here $\Delta_2(\kappa)$ and $\Delta_1(k)$ are matrices whose nonzero entries involve the differentials of the constituent complexes and the homomorphisms k and κ . It is clear that the dual complex is isomorphic to the complex obtained by taking $e^0\kappa$ and e^0k as representatives for k_D and $\kappa(D_*)$, respectively. \square

3. DUALITY FOR FINITE $\mathbb{F}\Lambda$ -MODULES

Let \mathbb{F} be a field and let $\mathbb{F}\Lambda = \mathbb{F}[t, t^{-1}]$. If M is a finite dimensional $\mathbb{F}\Lambda$ -module let $M^* = \overline{Hom_{\mathbb{F}\Lambda}(M, \mathbb{F}(t)/\mathbb{F}\Lambda)}$, $E(M) = \overline{Ext_{\mathbb{F}\Lambda}^1(M, \mathbb{F}\Lambda)}$ and $F(M) = Hom_{\mathbb{F}}(M, \mathbb{F})$. (The left Λ -module structure on the latter group is given by $(t\phi)(m) = \phi(t^{-1}m)$ for all $m \in M$ and $\phi : M \rightarrow \mathbb{F}$.) These modules are each non-canonically isomorphic to M , since $\mathbb{F}\Lambda$ is a PID. The functors $(-)^*$, $E(-)$ and $F(-)$ each define an involution on the category of finite dimensional $\mathbb{F}\Lambda$ -modules. They are in fact naturally equivalent. We shall give a simple proof of this for the case when \mathbb{F} is finite.

If $f \in \mathbb{F}\Lambda$ then $\text{Res}(f \frac{dt}{t}, \infty) + \text{Res}(f \frac{dt}{t}, 0) = 0$, since the only poles of $f \frac{dt}{t}$ are at 0 and ∞ . Thus we may define an \mathbb{F} -linear function $R : \mathbb{F}(t)/\mathbb{F}\Lambda \rightarrow \mathbb{F}$ by setting $R(q + \mathbb{F}\Lambda) = \text{Res}(q \frac{dt}{t}, \infty) + \text{Res}(q \frac{dt}{t}, 0)$ for all $q \in \mathbb{F}(t)$. In particular, if $q = \frac{f}{t^n - 1}$ where $f \in \mathbb{F}[t]$ has degree $< n$ then $R(q + \mathbb{F}\Lambda) = f(0)$.

Theorem 2. *There are natural equivalences $M^* \cong E(M) \cong F(M)$ on the category of finite dimensional $\mathbb{F}\Lambda$ -modules, if \mathbb{F} is a finite field.*

Proof. Let $0 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0$ be a free resolution of M . If $\phi \in M^*$ let $\phi_0 : P_0 \rightarrow \mathbb{F}(t)$ be a lift of ϕ . Then $\phi_0 \partial_1$ has image in $\mathbb{F}\Lambda$, and so defines a homomorphism $\phi_1 : P_1 \rightarrow \mathbb{F}\Lambda$ such that $\phi_1 \partial_2 = 0$. Consideration of the short exact sequence of complexes

$$0 \rightarrow Hom_{\mathbb{F}\Lambda}(P_*, \mathbb{F}\Lambda) \rightarrow Hom_{\mathbb{F}\Lambda}(P_*, \mathbb{F}(t)) \rightarrow Hom_{\mathbb{F}\Lambda}(P_*, \mathbb{F}(t)/\mathbb{F}\Lambda) \rightarrow 0$$

shows that $\delta_M(\phi) = [\phi_1]$, where $\delta_M : M^* \rightarrow E(M)$ is the Bockstein homomorphism associated to the coefficient sequence. (The extension corresponding to $\delta_M(\phi)$ is the pullback over ϕ of the sequence $0 \rightarrow \mathbb{F}\Lambda \rightarrow \mathbb{F}(t) \rightarrow \mathbb{F}(t)/\mathbb{F}\Lambda \rightarrow 0$.) Since M is a torsion $\mathbb{F}\Lambda$ -module and $\mathbb{F}(t)$ is divisible δ_M is an isomorphism.

Define $\chi_M : M^* \rightarrow F(M)$ by $\chi_M(\phi)(m) = R(\phi(m))$ for all $\phi \in M^*$ and $m \in M$. Then χ_M is $\mathbb{F}\Lambda$ -linear. Suppose now that M has finite order n . If $g \in F(M)$ let $L_M(g)(m) = \frac{\hat{g}(m)}{t^n - 1} + \mathbb{F}\Lambda$, where $\hat{g}(m) = \sum_{0 \leq k < n} g(t^k m) t^{-k}$ for all $m \in M$. Then $L_M(g)$ is $\mathbb{F}\Lambda$ -linear, since $(t^n - 1)M = 0$, and $\chi_M(L_M(g)) = g$. (This is the only point at which we need \mathbb{F} and M to be finite). Since M^* and $F(M)$ have the same dimension and χ_M is $\mathbb{F}\Lambda$ -linear χ_M and L_M are mutually inverse isomorphisms. \square

Let $\tau_M = \chi_M \delta_M^{-1} : E(M) \rightarrow F(M)$ be the composite isomorphism.

Theorem 2 holds also for infinite fields, but verifying that χ is an equivalence involves slightly more work. See [23] for the case $\mathbb{F} = \mathbb{Q}$.

There is also a corresponding result over Λ , i.e., with integer coefficients. Levine showed that if M is a finite Λ -module such that $(t^k - 1)M = 0$ then $e^2 M \cong \overline{Ext_{\Lambda}^1(M, \Lambda/(t^k - 1))} \cong \overline{Hom_{\Lambda}(M, \mathbb{Q}/\mathbb{Z} \otimes \Lambda/(t^k - 1))}$ (via appropriate Bockstein homomorphisms), which is in turn isomorphic to $Hom_{\mathbb{Z}}(M, \mathbb{Q}/\mathbb{Z})$ (via an explicit homomorphism similar to L_M) [17]. In particular, if M has exponent n as an

abelian group then $e^2M \cong \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z}/n\mathbb{Z})$. If A is a finitely generated Λ -module let $\sigma_A : e^2A \rightarrow \text{Hom}_{\mathbb{Z}}(T(A), \mathbb{Q}/\mathbb{Z})$ be the composite of these isomorphisms with the natural isomorphism $e^2A = e^2T(A)$.

4. ALGEBRAIC INVARIANTS FOR THE EXTERIOR OF A SIMPLE $2q$ -KNOT

Let K be a simple $2q$ -knot with $q > 2$. Then $\pi_1(X) = Z$ and $H_i(X; \Lambda) = 0$ for $0 < i < q$. Let $A = H_q(X; \Lambda)$ and $B = H_{q+1}(X; \Lambda)$. As observed in §1, these are torsion Λ -modules on which $t - 1$ acts invertibly. The Universal Coefficient spectral sequence and Poincaré duality give isomorphisms $e^1A = e^1(A/T(A)) \cong B$ and $e^2A \cong T(\overline{H^{q+2}(X; \Lambda)}) \cong T(A)$, while $H_i(X; \Lambda) = 0$ for $i > q + 1$. In particular, there is a nonsingular pairing $\ell : T(A) \times T(A) \rightarrow \mathbb{Q}/\mathbb{Z}$ and $p.d._{\Lambda} B \leq 1$. Since $B = (t - 1)B$ it follows that B is \mathbb{Z} -torsion free [17]. The exterior X fibres over S^1 if and only if A is finitely generated as an abelian group [3].

We may identify A with the Λ -module $\pi_q(X) = \pi_q(X')$, by the Hurewicz Theorem. Since $\pi_1(X) \cong Z$ and $\pi_j(X) = 0$ for $1 < j < q$ the Postnikov q -stage of X is the generalized Eilenberg-Mac Lane space $L_Z(A, q)$ determined by the Λ -module A . (See page 214 of [2].) Let $\tilde{\pi}_{q+1} = \pi_{q+1}(X)$, considered as a Λ -module, and let $\kappa(X) \in H^{q+2}(L_Z(A, q); \tilde{\pi}_{q+1})$ be the k -invariant for the next stage. Let $\eta \in \pi_3(S^2)$ be the Hopf map. Then $\eta_q = \Sigma^{q-2}\eta$ generates $\pi_{q+1}(S^q) = \pi_1^{st} = Z/2Z$ for all $q > 2$.

Since X' is $(q - 1)$ -connected there is an exact sequence

$$(2) \quad \pi_{q+2}(X') \rightarrow H_{q+2}(X'; \mathbb{Z}) \xrightarrow{b} A/2A \xrightarrow{\eta_q^*} \tilde{\pi}_{q+1} \rightarrow H_{q+1}(X'; \mathbb{Z}) \rightarrow 0,$$

where η_q^* is induced by composition with η_q and the unlabeled homomorphisms are Hurewicz homomorphisms. (See Theorem XII.3.12 of [25].) In our situation $H_{q+2}(X'; \mathbb{Z}) = 0$ and the right-hand portion of this sequence reduces to a short exact sequence of Λ -modules. Thus $\tilde{\pi}_{q+1}$ is determined by A and a class in $\text{Ext}_{\Lambda}^1(B, A/2A) = \text{Ext}_{\Lambda}^1(e^1(A/T(A)), A/2A)$. (See Theorem XII.4.16 of [25] for more on the corresponding class in $\text{Ext}_{\mathbb{Z}}^1(B, A/2A)$.) There is an analogous sequence when $q = 2$, involving the universal quadratic functor $\Gamma(A)$ instead of $A/2A$. In this case $\tilde{\pi}_{q+1} = \tilde{\pi}_3$ is not a stable homotopy group, there are nontrivial Whitehead products in the image of $\Gamma(A)$ and η has infinite order.

The chain complex of the covering space X' is naturally a complex of Λ -modules, and is chain homotopy equivalent to a finitely generated free complex C_* . As $H_j(C_*) = 0$ if $j > q + 2$ and $p.d._{\Lambda} H_{q+1}(C_*) \leq 1$ we may assume that C_* is the direct sum of the standard resolution of the augmentation Λ -module \mathbb{Z} and a simple Λ -complex C_*^{rel} concentrated in degrees $[q, q + 2]$. (This is homotopy equivalent to the equivariant chain complex of the pair (X, S^1) where S^1 is the meridian.) Therefore X is homotopy equivalent to a CW complex of dimension $q + 2$ [24]. Poincaré duality for $(X, \partial X)$ reduces to a chain homotopy equivalence $D(C_*^{rel})[-2q - 2]_* \simeq C_*^{rel}$, and so it follows from Theorem 1 that C_*^{rel} and C_* are determined up to chain homotopy equivalence by A .

The universal covering space of $L_Z(A, q)$ is $K(A, q)$, and the spectral sequence for the classifying map from $L_Z(A, q)$ to $S^1 = K(Z, 1)$ gives an exact sequence

$$(3) \quad 0 \rightarrow \text{Ext}_{\Lambda}^2(A, \tilde{\pi}_{q+1}) \rightarrow H^{q+2}(L_Z(A, q); \tilde{\pi}_{q+1}) \rightarrow \text{Hom}_{\Lambda}(A/2A, \tilde{\pi}_{q+1}) \rightarrow 0.$$

The image of $\kappa(X)$ in $\text{Hom}_{\Lambda}(A/2A, \tilde{\pi}_{q+1})$ is the homomorphism η_q^* . (It suffices to check the corresponding assertion for $\kappa(X')$, which follows from Lemma 4.15 of

Chapter XII of [25], since $\text{Hom}_\Lambda(A/2A, \tilde{\pi}_{q+1}) \leq \text{Hom}_\mathbb{Z}(A/2A, \tilde{\pi}_{q+1})$.) The corresponding sequence with coefficients B reduces to an isomorphism $\text{Ext}_\Lambda^2(A, B) \cong H^{q+2}(L_Z(A, q); B)$, since B is \mathbb{Z} -torsion free, and the image of $\kappa(X)$ under the coefficient homomorphism induced by h is $\kappa(C_*)$.

Theorem 3. *Let A and $\tilde{\pi}_{q+1}$ be finitely generated Λ -torsion modules and $\kappa \in H^{q+2}(L_Z(A, q); \tilde{\pi}_{q+1})$. Suppose that the image of κ in $\text{Hom}_\Lambda(A/2A, \tilde{\pi}_{q+1})$ is a monomorphism with cokernel B , and that $p.d._\Lambda B \leq 1$. Then there is a finite $(q+2)$ -complex X_κ such that $\pi_1(X_\kappa) \cong Z$, $\pi_q(X_\kappa) \cong A$, $\pi_{q+1}(X_\kappa) \cong \tilde{\pi}_{q+1}$, $\tilde{H}_j(X_\kappa; \Lambda) = 0$ if $j \neq q$ or $q+1$, and $\kappa(X) = \kappa$. The homotopy type of X is uniquely determined by A , $\tilde{\pi}_{q+1}$ and κ .*

Proof. These invariants determine a Postnikov $(q+1)$ -stage $P = P_{q+1}(\kappa)$. We may assume the k -skeleton $P^{[k]}$ is finite for all $k \geq 0$, since A and $\tilde{\pi}_{q+1}$ are finitely generated Λ -modules and Λ is noetherian. The hypothesis on κ implies that $H_{q+1}(P^{[q+2]}; \Lambda) = H_{q+1}(P; \Lambda) \cong B$ and that composition with η_q determines a monomorphism from $A/2A$ to $\tilde{\pi}_{q+1}$. Hence the Hurewicz homomorphism for $P^{[q+2]}$ in degree $q+2$ is onto. Since $p.d._\Lambda B \leq 1$ and $P^{[q+2]}$ is a finite $(q+2)$ -complex $H_{q+2}(P^{[q+2]}; \Lambda)$ is a finitely generated free module. Attaching $(q+3)$ -cells to $P^{[q+2]}$ along representatives for a basis of $H_{q+2}(P^{[q+2]}; \Lambda)$ gives a finite $(q+3)$ -complex with trivial homology in degrees $\geq q+2$, and which is therefore homotopy equivalent to a finite $(q+2)$ -complex X_κ , by [24].

Let Y be a $(q+2)$ -complex with Postnikov $(q+1)$ -stage $f : Y \rightarrow P$ and such that $H_{q+2}(Y; \Lambda) = 0$. Then f factors through a map $g : Y \rightarrow X_\kappa$, since we may construct P by adjoining cells of dimension $\geq q+3$ to X_κ , to kill the higher homotopy groups. The lift of g to the universal covers induces isomorphisms on homology and so is a homotopy equivalence, by the Hurewicz and Whitehead Theorems. Therefore g is a homotopy equivalence. \square

It is easy to construct a finite $(q+2)$ -complex with equivariant chain complex of the required chain homotopy type, using only the surjectivity of the Hurewicz homomorphism in degree $q+1$ (and not appealing to [24]).

However not all such complexes admit Poincaré duality isomorphisms. Let $D_* = D(C^{rel})[-2q-2]_*$. It is clear from Theorem 1 that there is a chain homotopy equivalence $D_* \simeq C_*^{rel}$ if and only if $k(A)$ and $\kappa(C_*)$ correspond.

An analogous but simpler argument shows that if K is a simple $(2q-1)$ -knot the homotopy type of $X = X(K)$ is determined by the module $A = H_q(X; \Lambda)$ alone.

5. FARBER QUINTUPLES AND KEARTON F -FORMS

The Farber quintuple of K is $q_F(K) = (A, \Pi, \alpha, \ell, \psi)$, where $A = H_q(X; \Lambda)$, $\Pi = \pi_{q+2}^{st}(X')$, $\alpha : A/2A \rightarrow \Pi$ is the monomorphism determined by composition with $\eta_q \eta_{q+1}$, $\ell : T(A) \times T(A) \rightarrow \mathbb{Q}/\mathbb{Z}$ is the torsion linking pairing and $\psi : \Pi \times \Pi \rightarrow \mathbb{Z}/4\mathbb{Z} < \pi_3^{st}$ is a bilinear pairing derived from Spanier-Whitehead duality for a Seifert hypersurface for K . The pairings ℓ and ψ are $(-1)^{q+1}$ -symmetric, nonsingular and t acts isometrically. In other words the *adjoint* functions $Ad(\ell)$ and $Ad(\psi)$ are Λ -isomorphisms, where $Ad(\ell) : T(A) \rightarrow \text{Hom}_\mathbb{Z}(T(A), \mathbb{Q}/\mathbb{Z})$ is given by $Ad(\ell)(b)(a) = \ell(a, b)$ for all $a, b \in T(A)$, and $Ad(\psi)$ is defined similarly. Let $\beta = \alpha^* Ad(\psi) : \Pi \rightarrow \text{Hom}_{\mathbb{F}_2}(A/2A, \mathbb{F}_2)$, so that $\beta(p)(a) = \psi(p, \alpha(a))$ for all $a \in A/2A$ and $p \in \Pi$. Then β is an epimorphism and $\text{Ker}(\beta) = \text{Im}(\alpha)$.

These pairings interact in the following way. If M is a Λ -module and $m \in M$ let $[m]_2$ be the image of m in $M/2M$. Let $\gamma : \Pi \rightarrow T(A)$ be the homomorphism determined by $\psi(p, \alpha([x]_2)) = \ell(\gamma(p), x)$ for all $x \in T(A)$ and $p \in \Pi$ and nonsingularity of ℓ . Then $\alpha([\gamma(p)]_2) = 2p$, for all $p \in \Pi$. (In particular, $\gamma = 0$ if and only if $T(A)$ has odd order, and then Π has exponent 2, ℓ and ψ are independent and ψ is symmetric.) There is an obvious notion of sum of Farber quintuples, and $q_F(K \# \tilde{K}) = q_F(K) \oplus q_F(\tilde{K})$.

In the torsion free case ($T(A) = 0$) there is an alternative system of invariants, due to Kearton [14]. Reducing sequence (2) above modulo (2) gives a sequence

$$\mathcal{E} : 0 \rightarrow A/2A \rightarrow \tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1} \rightarrow B/2B \rightarrow 0$$

which is exact since B is \mathbb{Z} -torsion free. The Kearton F -form of K involves the exact sequence \mathcal{E} and an hermitean pairing $[-, -]$ on $\tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$ with values in $\mathbb{F}_2(t)/\mathbb{F}_2\Lambda$ (and some auxiliary data). The pairing is defined geometrically, in terms of intersection numbers. It may be identified with an isometric structure on a \mathbb{F}_2 -valued pairing, via the natural isomorphism $\tau_M : M^* \rightarrow F(M)$ defined in §2 above, and the F -form may be reformulated as a pair (\mathcal{E}, b) , where $b : \mathcal{E} \rightarrow \mathcal{E}^*$ is an isomorphism such that $b^* = (-1)^qb$ [26]. It is a complete invariant for torsion free simple $2q$ -knots if $q \geq 3$ [9]. (The pairing $[-, -]$ is defined for any simple $2q$ -knot, and is nonsingular if $T(A)$ is odd. The latter hypothesis is needed only for Lemmas 1.10 and 1.13 of §1 of [14].)

These invariants are defined for $q \geq 1$, but simple 2-knots are topologically trivial, and the natural invariants for simple 4-knots are not stable, so it is unlikely that the Farber quintuple is a complete invariant for these. (See instead [11].) In the torsion free case $\Pi \cong \tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$ and the Farber quintuple is algebraically equivalent to a Kearton F -form (although it is not clear how the pairings ψ and $[-, -]$ correspond). This suggests that the Farber quintuple may suffice when $q = 3$. Can the Farber invariant be extended to all even-dimensional knots as an additive invariant?

6. RELATING THE INVARIANTS

The first three ingredients of $q_F(K)$ are determined by the homotopy type of X alone, while ℓ and ψ are manifestations of Poincaré duality for the pair $(X, \partial X)$. We shall show that, conversely, $q_F(K)$ determines $\tilde{\pi}_{q+1}$ and $\kappa(X)$ and hence the homotopy type of X , at least if $T(A)$ has odd order. We shall consider how the duality pairings and the inclusion of ∂X interact in the next section.

Let $pd_Z : B = H_{q+1}(X; \Lambda) \rightarrow e^1 A$ and $pd_2 : H_{q+1}(X; \mathbb{F}_2\Lambda) \rightarrow E(A/2A) = \overline{Ext}_{\mathbb{F}_2\Lambda}^1(A/2A, \mathbb{F}_2\Lambda)$ be the isomorphisms determined by Poincaré duality and the Universal Coefficient spectral sequence, and let $\tau : E(A/2A) \cong Hom_{\mathbb{F}_2}(A/2A, \mathbb{F}_2)$ be the isomorphism defined in §2. Let $\rho_2 : B = H_{q+1}(X; \Lambda) \rightarrow H_{q+1}(X; \mathbb{F}_2\Lambda)$ be reduction *mod* (2). Reduction *mod* (2) also induces a natural isomorphism $\mathbb{F}_2 \otimes_{\mathbb{Z}} e^1(A/T(A)) \cong E(A/(T(A), 2A))$ and hence a natural transformation $e : e^1 A \cong e^1(A/T(A)) \rightarrow E(A/(T(A), 2A)) \rightarrow E(A/2A)$, and the diagram

$$(4) \quad \begin{array}{ccccc} B & \longrightarrow & H^{q+1}(X; \Lambda) & \longrightarrow & e^1 A \\ \downarrow \rho_2 & & \downarrow & & \downarrow e \\ H_{q+1}(X; \mathbb{F}_2\Lambda) & \longrightarrow & H^{q+1}(X; \mathbb{F}_2\Lambda) & \longrightarrow & E(A/2A) \end{array}$$

is commutative.

Theorem 4. *The module $\tilde{\pi}_{q+1}$ is the pullback of τe and β over $\text{Hom}_{\mathbb{F}_2}(A/2A, \mathbb{F}_2)$, i.e., it is the fibre sum $\tilde{\pi}_{q+1} \cong \{(b, p) \in e^1 A \times \Pi \mid \tau e(b) = \beta(p)\}$.*

Proof. The Atiyah-Hirzebruch spectral sequence for the stable homotopy of X' gives the exact sequence (2) (if $q > 2$) and also an exact sequence

$$(5) \quad 0 \rightarrow H_q(X'; \pi_2^{st}) \rightarrow \Pi \xrightarrow{\theta} H_{q+1}(X'; \pi_1^{st}) \rightarrow 0.$$

We may identify $H_q(X'; \pi_1^{st})$ and $H_q(X'; \pi_2^{st})$ with $A/2A$, since $\pi_1^{st} = \pi_2^{st} = Z/2Z$. This sequence is an exact sequence of Λ -modules, by the naturality of the Atiyah-Hirzebruch spectral sequence. Moreover $\beta = \tau p d_2 \theta$. (See Theorem 3.3(b) of [6].) Composition (on the right) with η_{q+1} induces a homomorphism $\eta_{q+1}^* : \tilde{\pi}_{q+1} \rightarrow \Pi$, which factors through $\tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$, since η_{q+1} has order 2. The sequences (2) and (5) form the rows of a commutative diagram

$$(6) \quad \begin{array}{ccccccc} 0 & \longrightarrow & A/2A & \xrightarrow{\eta_q^*} & \tilde{\pi}_{q+1} & \longrightarrow & B = H_{q+1}(X'; \mathbb{Z}) \longrightarrow 0 \\ \downarrow & & \downarrow = & & \downarrow \eta_{q+1}^* & & \downarrow \rho_2 & \downarrow \\ 0 & \longrightarrow & A/2A & \xrightarrow{\alpha} & \Pi & \xrightarrow{\theta} & H_{q+1}(X'; \mathbb{F}_2) & \longrightarrow 0. \end{array}$$

The theorem now follows easily. \square

Reduction *mod* (2) induces a monomorphism from $B/2B$ to $H_{q+1}(X'; \mathbb{F}_2)$, with cokernel $\text{Tor}(A, Z/2Z)$, while η_{q+1}^* factors through $\tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$, since $2\eta_{q+1} = 0$.

Corollary 5. *If $T(A)$ has odd order the homotopy type of X is determined directly by $q_F(K)$.*

Proof. Since $T(A)$ has odd order $\text{Tor}(A, Z/2Z) = 0$, so η^* is an epimorphism, and $\text{Ext}_{\Lambda}^2(A, \tilde{\pi}_{q+1}) \cong \text{Ext}_{\Lambda}^2(T(A), e^1(A/T(A)))$, and is the odd-order summand of $H^{q+2}(L_Z(A, q); \tilde{\pi}_{q+1})$.

Since the image of $\kappa(X)$ in the 2-primary summand is η_q^* it follows that $[\kappa(X)]$ is determined by $[\kappa(C)]$ and α , and hence by A and α . Thus the homotopy type of X is determined directly by $q_F(K)$, as in Theorems 3 and 4. \square

If $A = 2A$ then $\Pi = 0$ and so $q_F(K)$ reduces to the pair (A, ℓ) . In this case $\tilde{\pi}_{q+1} \cong e^1 A$ and $\kappa(X) = \kappa(C) \in \text{Ext}_{\Lambda}^2(A, e^1 A)$. If instead A is torsion free $\text{Ext}_{\Lambda}^2(A, \tilde{\pi}_{q+1}) = 0$ and so $\kappa(C) = 0$. Hence $\kappa(X)$ is determined by α , which lifts to a monomorphism $A/2A \cong \{(0, p) \in B \times \Pi \mid \beta(p) = 0\} \leq \tilde{\pi}_{q+1}$ in the above fibre sum. (In this case $\kappa(C) = 0$.)

We may also ask how we may derive Π and α directly from A , $\tilde{\pi}_{q+1}$ and $\kappa(X)$. If $T(A)$ is odd $\Pi \cong \tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$ and α is the composite of the monomorphism η_q^* with reduction *mod* (2). In this case $q_F(K)$ is algebraically equivalent to the torsion linking pairing ℓ together with an F -form (with associated module $A/T(A)$).

In [6] the pairing ψ is defined in terms of Spanier-Whitehead duality for a Seifert hypersurface V for the knot K . Let $u : V \wedge V \rightarrow S^{n+1}$ and $z : V \rightarrow V$ be the duality pairing and carving map of [6], respectively. Let $\psi_V(\alpha, \beta) = u(\alpha \wedge \beta) \in \pi_{2q+4}^{st}(S^{2q+1}) = \pi_3^{st} \cong Z/24Z$ for all $\alpha, \beta \in \pi_{q+2}^{st}(V)$. Let $P = \mathbb{Z}[z]$, $\bar{z} = 1 - z$ and $L = P[(z\bar{z})^{-1}]$. Then $\pi_{q+2}^{st}(X') \cong L \otimes_P \pi_{q+2}^{st}(V)$ and ψ_V extends to a pairing ψ on $\pi_{q+2}^{st}(X')$ which is independent of the choice of Seifert hypersurface V . Is there a direct definition of ψ which does not involve the choice of a Seifert hypersurface?

For instance, can it be defined directly in terms of a Z -equivariant SW duality? (See also [5] for the case of fibred simple $2q$ -knots.)

We may define similar pairings Ψ_V and Ψ on $\pi_{q+1}^{st}(V)$ and $\pi_{q+1}^{st}(X')$, with values in $\pi_1^{st} = Z/2Z$. Composition on the right with η^2 induces a monomorphism from π_1^{st} to π_3^{st} , and $\eta \wedge \eta = \eta^2 \in \pi_2^{st}$, by X.8.12 of [25]. Hence $\Psi(a, b) = \psi(a\eta_{q+1}, b\eta_{q+1})$ for all $a, b \in \pi_{q+1}^{st}(X')$. If $T(A)$ is odd η_{q+1}^* induces an isomorphism $\pi_{q+1}^{st}(X') \cong \pi_{q+2}^{st}(X')$ and so ψ and Ψ are equivalent. We expect that the pairings $[-, -]$ and Ψ may then be related as follows:

Lemma? *If $T(A)$ is odd $\Psi(u, v) = R([u, v])$ for all $u, v \in \tilde{\pi}_{q+1}/2\tilde{\pi}_{q+1}$.*

7. INCLUSION OF THE BOUNDARY

Let $Y = S^{2q} \times S^1$ have the product orientation and let $J_K : Y \cong \partial X \subset X$ be the inclusion of the boundary. The oriented homotopy type of X and the class $[J(K)]$ of J_K modulo homotopy and composition with orientation preserving self homotopy equivalences of X and Y together determine the oriented homotopy type of the pair $(X, \partial X)$. The difficulty is in giving a practical characterization of $[J(K)]$. In this section we shall mention two of the known constraints.

Let $J : Y \rightarrow X$ be a map which induces an isomorphism on fundamental groups. Let $M(J)$ be the mapping cylinder of J . There is a canonical homotopy equivalence $M(J) \simeq X$. The orientation for Y determines a generator μ_J for $H_{2q+2}(M(J), Y; \mathbb{Z}) \cong H_{2q+1}(Y; \mathbb{Z})$. Since the universal covers of $M(J) \simeq X$ and Y are highly connected cap product with μ_J determines homomorphisms $\overline{H^{q+1}(X; \Lambda)} \rightarrow H_{q+1}(X; \Lambda)$ and $\overline{H^{q+2}(X; \Lambda)} \rightarrow H_q(X; \Lambda)$. The pair $(M(J), Y)$ is a PD_{2q+2} -pair if and only if these homomorphisms are isomorphisms. (The corresponding homomorphisms in other degrees are isomorphisms, in all cases.)

The map $J' : Y' = S^{2q} \times R \rightarrow X'$ covering J determines an element $\lambda_J \in \pi_{2q}(X') = \pi_{2q}(X)$, such that $(t-1)\lambda_J = 0$. In the fibred case the Poincaré duality constraints depend only on λ_J . The longitude λ_K corresponds to J_K . Lemma 3 of [11] asserted that *if X is an n -dimensional homology circle with $H_n(X; \Lambda) = 0$ and two maps $j_1, j_2 : S^n \times S^1 \rightarrow X$ induce isomorphisms on π_1 and agree on $S^n \vee S^1$ then there is a self-homotopy equivalent $f : X \rightarrow X$ such that $j_2 \sim fj_1$* . If this were correct it would follow not only that λ_K determines J_K for 1-simple knots but also that such knots are determined by their exteriors. The proof of Lemma 3 in [11] is wrong, and thus the later assumptions in that paper that the homotopy quadruple discussed there is a complete invariant for simple 4-knots are unjustified. (We thank W.Richter for pointing out our error). Nevertheless it remains possible that the result may be true, at least if X' is sufficiently highly connected.

The inclusion of the boundary of a Seifert hypersurface V for K has trivial suspension in $\pi_{2q+1}(SV)$ [22]. Therefore λ_K is in the kernel of the suspension $E : \pi_{2q}(X') \rightarrow \pi_{2q+1}(SX')$. The group $\pi_{2q}(X')$ is just outside the stable range, since X' is $(q-1)$ -connected, and sits near one extremity of the EHP sequence (Theorem XII.2.2 of [25]):

$$\pi_{3q-2}(X') \rightarrow \cdots \rightarrow \pi_{2q+2}(SX) \rightarrow \pi_{2q+2}(X' * X') \rightarrow \pi_{2q}(X') \rightarrow \pi_{2q+1}(SX') \rightarrow 0.$$

Moreover $X' \simeq M_A \vee M_B$ is a wedge of Moore spaces $M_A = M(A, q)$ and $M_B = M(B, q+1)$, since X' is $(q-1)$ -connected, $H_j(X'; \mathbb{Z}) = 0$ for $j > q+1$ and $q > 2$. Hence $\pi_{2q}(X') \cong \pi_{2q}(M_A) \oplus \pi_{2q}(M_B) \oplus [A, B]$, where $[A, B]$ is the subgroup generated by all Whitehead products $[x, y]$ with $x \in \pi_q(M_A) \cong A$ and

$y \in \pi_{q+1}(M_B) \cong B$. The summand $\pi_{2q}(M_B)$ is stable, while all Whitehead products have trivial suspension.

Suppose now that K is fibred and $A \cong Z^\beta$ is torsion-free. Then $M_A \simeq \vee^\beta S^q$ and $M_B \simeq \vee^\beta S^{q+1}$. Let $\{i_q^m\}_{m \leq \beta}$ and $\{i_{q+1}^n\}_{n \leq \beta}$ be bases for A and the image of $B \cong \pi_{q+1}(\vee^\beta S^{q+1})$ in $\pi_{q+1}(X')$, respectively. In this case $\text{Ker}(E)$ is generated by all Whitehead products $i_{mm'} = [i_q^m, i_q^{m'} \eta_{q+1}]$ and $j_{mn} = [i_q^m, i_{q+1}^n]$ for all $m, m', n \leq \beta$. Thus $\lambda_K = \sum c_{mm'} i_{mm'} + \sum d_{mn} j_{mn}$ for some coefficients $c_{mm'} \in \mathbb{F}_2$ and $d_{mn} \in \mathbb{Z}$. The integral coefficients d_{mn} may be detected by computing cup products in $(S^q \vee S^{q+1}) \cup_{p\lambda} e^{2q+1}$ for suitable projections $p : X' \rightarrow S^q \vee S^{q+1}$. (See [12], and note that $S^q \vee S^{q+1} \cup_{[i,j]} e^{2q+1} \simeq S^q \times S^{q+1}$.) Hence the matrix $D = [d_{mn}]$ is invertible, since (X', λ_K) is a PD_{2q+1} -complex. and we may choose the basis for B so that $D = I_\beta$.

Question. *How are the coefficients $c_{mm'}$ determined by the pairing Ψ ?*

An analogous but simpler argument applies if K is a fibred simple $(4k-1)$ -knot. For then $X' \simeq \vee^\beta S^{2k}$ and $\text{Ker}(E) \leq \pi_{4k-1}(X')$ is generated by the Whitehead products $[i_{2k}^m, i_{2k}^n]$ for $m, n \leq \beta$. The coefficients e_{mn} of $\lambda_K = \sum e_{mn} [i_{2k}^m, i_{2k}^n]$ may be detected by cup products in $S^{2k} \cup_{p\lambda} e^{4k}$ or $(S^{2k} \vee S^{2k}) \cup_{p\lambda} e^{4k}$ for suitable projections $p : X' \rightarrow S^{2k}$ (and using the Hopf invariant, as on page 211 of [18]) or $p : X' \rightarrow S^{2k} \vee S^{2k}$ (and using [12]).

When $T(A) \neq 0$ there may be elements of $\text{Ker}(E)$ which are not Whitehead products, as can be seen already when $A = Z/pZ$. This is the situation that arises for *odd finite* simple $2q$ -knots, for which $A = T(A)$ is finite of odd order and the Farber quintuple reduces to the torsion linking pairing ℓ . (See [15].)

8. HERMITIAN SELF-DUALITIES

Let \mathcal{C} be an additive category with an involution $*$ and let $\varepsilon = \pm 1$. An ε -hermitean pairing on an object C of \mathcal{C} is defined to be an isomorphism $\phi : C \rightarrow C^*$ such that $\phi^* = \varepsilon\phi$. This categorical point of view is particularly useful if the Krull-Schmidt Theorem holds for the objects of \mathcal{C} and the endomorphism rings of objects are radically complete. For then the analysis of the decompositions of hermitean pairings into orthogonal direct sums may be reduced to corresponding questions for pairings over (skew) fields [19].

Consider quadruples $Q = (A, C, \theta, \mathcal{E})$ where A and C are finitely generated Λ -modules on which $t-1$ acts invertibly, $\theta : T(A) \rightarrow e^2 C$ is an isomorphism and \mathcal{E} is an exact sequence of Λ -modules

$$(7) \quad 0 \rightarrow A/2A \xrightarrow{\alpha} \Pi \xrightarrow{\omega} e^2(C/2C) \rightarrow 0.$$

A morphism between two such quadruples is a triple $\phi = (f, g, h)$ where $f : A \rightarrow A'$, $g : C' \rightarrow C$ and $h : \Pi \rightarrow \Pi'$ such that $\theta' f|_{T(A)} = e^2 g \theta$, $h \alpha = \alpha' [f]_2$ and $\omega' h = e^2 [g]_2 \omega$. The category \mathcal{Q} of such quadruples is additive.

Define a duality functor by $Q^* = (A, C, \theta, \mathcal{E})^* = (C, A, e^2 \theta, e^2 \mathcal{E})$ and $(f, g, h)^* = (g, f, e^2 h)$. An isomorphism $\phi = (f, g, h)$ from Q to Q^* is an ε -symmetric self-duality of Q if $\phi^* = \varepsilon\phi$, in which case f and g are isomorphisms from A to C , $f = \varepsilon g$ and $h = \varepsilon e^2 h$. A pair (Q, ϕ) with ϕ an ε -symmetric self-duality of q is *hyperbolic* if there is an object N in \mathcal{Q} such that $Q \cong N \oplus N^*$ and ϕ has matrix $\begin{pmatrix} 0 & \varepsilon \\ 1 & 0 \end{pmatrix}$ with respect to this decomposition.

Theorem 6. *Farber quintuples for $2q$ -knots correspond bijectively to pairs (Q, ϕ) where $Q = (A, C, \theta, \mathcal{E})$ is an object in \mathcal{Q} and $\phi = (f, g, h)$ is a $(-1)^{q+1}$ -symmetric self-duality of Q such that $\alpha[\gamma(p)]_2 = 2p$ for all $p \in \Pi$, where $\gamma(p)$ is determined by $e^2 g\theta(\gamma(p)(x)) = \sigma_\Pi h(\alpha[x]_2)(p)$ for all $x \in T(A)$ and $p \in \Pi$.*

Proof. Let $(A, \Pi, \alpha, \ell, \psi)$ be a Farber quintuple with $(-1)^{q+1}$ -symmetric pairings ℓ and ψ and let β be the associated epimorphism. Let $C = A$, $\theta = \sigma_A^{-1} \text{Ad}(\ell)$, $\omega = \sigma_A^{-1} \beta$ and \mathcal{E} be the exact sequence determined by α and ω . Then $Q = (A, A, \theta, \mathcal{E})$ is in \mathcal{Q} and $(1_A, 1_A, \sigma_\Pi^{-1} \text{Ad}(\psi))$ is a $(-1)^{q+1}$ -symmetric self-duality of Q .

Conversely, if $\phi = (f, g, h)$ is a $(-1)^{q+1}$ -symmetric self-duality of $(A, C, \theta, \mathcal{E})$ we may define pairings ℓ on $T(A)$ and ψ on Π by setting $\ell(a, a') = e^2 g\theta(a')(a)$ for all $a, a' \in T(A)$ and $\psi(p, q) = \sigma_\Pi(h(q))(p)$ for all $p, q \in \Pi$. (We then have $\beta = \sigma_A e^2 g\omega$.) Then $(A, \Pi, \alpha, \ell, \psi)$ is almost a Farber quintuple; we need only the additional condition that $\alpha[\gamma(p)]_2 = 2p$ for all $p \in \Pi$, where γ is defined as in §5. The equation in the statement of the theorem is a reformulation of this condition in terms of the constituents of ϕ . \square

The Krull-Schmidt Theorem holds in any additive category in which (i) all idempotents split; and (ii) every object is a finite sum of objects whose endomorphism rings are local rings. (Such summands are necessarily indecomposable). It is easily verified that \mathcal{Q} satisfies the first condition. We shall determine the objects of \mathcal{Q} which satisfy (ii) by means of the next lemma.

Lemma 7. *Let R be a local Λ -algebra which is finitely generated as a Λ -module. Then R is finite.*

Proof. Let $p : \Lambda \rightarrow R$ be the natural homomorphism. Let $J = \text{rad}(R) = R - R^\times$ and $m = p^{-1}(J)$. The skewfield R/J is finitely generated as a Λ/m -module. It follows easily that Λ/m is a field. In particular m is a maximal ideal (and R/J is a finite field). Since $p(\Lambda)_m \leq R$ it is an integral extension of $p(\Lambda)$, and every prime ideal of $p(\Lambda)$ is the restriction of a prime ideal of $p(\Lambda)_m$, by Theorem 5.10 of [1]. Thus $p(\Lambda)$ is also a local ring, and so m is the unique maximal ideal of Λ which contains $K = \text{Ker}(p)$. Consideration of the primary decomposition of K shows that K must be m -primary and $p(\Lambda) = \Lambda/K$ finite. Hence R is also finite. \square

An object N in an abelian category \mathcal{C} is a *torsion* object if the ring $\text{End}_{\mathcal{C}}(N) = \mathcal{C}(N, N)$ is finite.

Theorem 8. *Let $Q = (A, C, \theta, \mathcal{E})$ in \mathcal{Q} . Then the following are equivalent*

- (1) *A and C are finite Λ -modules;*
- (2) *Q is a torsion object;*
- (3) *Q is a finite sum of objects whose endomorphism rings are local rings.*

The Krull-Schmidt Theorem holds in the full subcategory of \mathcal{Q} determined by the torsion objects, and the endomorphism rings of such objects are radically complete.

Proof. It is immediate that $\text{End}_{\mathcal{Q}}(Q)$ has finite index in $\text{End}_{\Lambda}(A) \times \text{End}_{\Lambda}(C)$. Thus $\text{End}_{\mathcal{Q}}(Q)$ is finite if and only if $\text{End}_{\Lambda}(A)$ and $\text{End}_{\Lambda}(C)$ are both finite. It follows easily that Q is a torsion object in \mathcal{Q} if and only if A and C are finite. Thus (1) \Leftrightarrow (2).

In a finite ring some power of each element is idempotent. Thus a finite ring with no nontrivial idempotents is local. Since idempotents in \mathcal{Q} split it follows that (2) \Rightarrow (3).

If $\text{End}_{\mathcal{Q}}(Q)$ is local then it is finite, by the lemma, and so Q is a torsion object. Thus (3) \Rightarrow (2).

The final assertion is now clear, since the radical of an Artinian ring is nilpotent. \square

If $(A, C, \theta, \mathcal{E})$ is a torsion object in \mathcal{Q} then $\theta : A \cong e^2 C$ and $e^2(C/2C) \cong {}_2A$, and so the full subcategory of \mathcal{Q} determined by the torsion objects is essentially the category considered in [8].

9. DOUBLE NULL CONCORDANCE

We expect that the following assertions should be true.

Theorem? *A simple $2q$ -knot K with $q \geq 4$ is doubly null concordant if and only if $q_F(K)$ is hyperbolic.*

The necessity of the condition should be easy.

Corollary? *A simple $2q$ -knot K with $q \geq 4$ is stably DNC if and only if the Witt class of $q_F(K)$ is trivial.*

The sufficiency of the condition would follow from the theorem and the fact that every Farber quintuple is realized by some simple $2q$ -knot. The necessity of the condition may require extending $q_F(-)$ to an invariant of all even-dimensional knots.

REFERENCES

- [1] Atiyah, M.F. and Macdonald, I.G. *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, Mass. - London - Don Mills, Ont. (1969).
- [2] Baues, H.-J. *Algebraic Homotopy*, Cambridge UP, Cambridge (1989).
- [3] Browder, W. and Levine, J.P. Fiberings manifolds over a circle, *Comment. Math. Helvetici* 40 (1966), 153-160.
- [4] Dold, A. Zur Homotopietheorie der Kettenkomplexe, *Math. Ann.* 140 (1960), 278-298.
- [5] Farber, M.S. Classification of stable fibred knots, *Mat. Sbornik* 115 (1981), 223-262.
- [6] Farber, M.S. An algebraic classification of some even-dimensional spherical knots I, II, *Trans. Amer. Math. Soc.* 281 (1984), 507-528; 529-570.
- [7] Farber, M.S. and Hillman, J.A. Doubly null concordant knots have hyperbolic stable isometry structures, *J. Knot Theory Ramif.* 2 (1993), 125-140.
- [8] Hillman, J.A. Finite simple even-dimensional knots, *J. London Math. Soc.* 34 (1986), 369-374.
- [9] Hillman, J.A. and Kearton, C. Seifert matrices and 6-knots, *Trans. Amer. Math. Soc.* 309 (1988), 843-857.
- [10] Hillman, J.A. and Kearton, C. Stable isometry structures and the factorization of Q -acyclic stable knots, *J. London Math. Soc.* 39 (1989), 175-182.
- [11] Hillman, J.A. and Kearton, C. Simple 4-knots, *J. Knot Theory Ramif.* 7 (1998), 907-923.
- [12] James, I.M. Note on cup-products, *Proc. Amer. Math. Soc.* 8 (1957), 374-383.
- [13] Kearton, C. An algebraic classification of some even-dimensional knots, *Topology* 15 (1976), 363-373.
- [14] Kearton, C. An algebraic classification of certain simple even-dimensional knots, *Trans. Amer. Math. Soc.* 176 (1983), 1-53.
- [15] Kojima, S. Classification of simple knots by Levine pairings, *Comment. Math. Helvetici* 54 (1979), 356-367.
- [16] Lashof, R. and Shaneson, J.L. Classification of knots in codimension two, *Bull. Amer. Math. Soc.* 75 (1969), 171-175.
- [17] Levine, J.P. Knot modules I, *Trans. Amer. Math. Soc.* 229 (1977), 1-50.
- [18] May, J.P. *A Concise Course on Algebraic Topology*, Chicago University Press, Chicago (1999).
- [19] Quebbemann, H.-G., Scharlau, W. and Schulte, M. Quadratic and hermitean forms in additive categories, *J. Algebra* 59 (1979), 264-289.
- [20] Richter, W. High-dimensional knots with $\pi_1 \cong \mathbb{Z}$ are determined by their complements in one more dimension than Farber's range, *Proc. Amer. Math. Soc.* 120 (1994), 285-294.
- [21] Richter, W. A homotopy-theoretic proof of William's metastable Poincaré embedding theorem, *Duke Math. J.* 88 (1997), 535-447.
- [22] Suciuc, A.I. Inequivalent frame-spun knots with the same complement, *Comment. Math. Helvetici* 67 (1992), 47-63.
- [23] Trotter, H.S. Knot modules and Seifert matrices, in *Knot Theory - Proceedings, Plans sur Bex, Switzerland 1977* (edited by J.-C. Hausmann), Lecture Notes in Mathematics 685, Springer Verlag, Berlin - Heidelberg - New York (1978), 291-299.
- [24] Wall, C.T.C. Finiteness conditions for CW-complexes I, *Ann. Math.* 81 (1965), 56-69.
- [25] Whitehead, G.W. *Elements of Homotopy Theory*, Graduate Texts in Mathematics vol. 61, Springer-Verlag, Berlin - Heidelberg - New York (1978).
- [26] Wilson, S.M.J. Invariants of even-dimensional torsion-free knots, preprint, Durham (1989).

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA
E-mail address: john@maths.usyd.edu.au